



Seizures After Traumatic Brain Injury (TBI)

Days after getting a TBI in a car crash, Seneca had a seizure. It happened during rehab at an American Indian Health Center.



One out of 10 people who are hospitalized after a TBI will have a seizure.

Most seizures happen in the first few days or weeks after a TBI.

But some may occur months or even years after a TBI.



Before the seizure, Seneca had some signs. He saw flashing colors and felt tired all of a sudden.

What are these flashing colors?

Along with changes in vision and sudden tiredness, other signs include:

- **Unusual movement of the head, body, arms, legs, or eyes.**
- **Unresponsiveness and staring.**
- **Chewing, lip smacking, or fumbling movements.**
- **Sudden experience of an odd smell or taste.**
- **Not being able to speak or understand others.**
- **New outbursts of anger or tearfulness.**

And why do I feel tired all of a sudden?





Because Seneca is at risk for seizures in the future, Dr. Sky tells him the things that can increase the risk of another one.




Dr. Sky tells Seneca that he can take anti-seizure medicine to help control his seizures. Most people adjust well to their medication.

If a medication is causing persistent side-effects, the physician may be able to prescribe another medication.





Dr. Sky tells Seneca there are safety issues that he needs to be aware of if he keeps having seizures.



Dr. Sky also tells Seneca to make a diary about his seizures. Include the date, time of day, length of time, and a description of each seizure. Your doctor will need this information, along with the medicines you take to control the seizures. Most seizures are short and don't cause serious injuries.



**Thanks for
all the help,
Dr. Sky!**

**You are most welcome.
I also have this pocket guide
for you, which covers what
caregivers should do for
someone having a seizure.**



A Pocket Guide for Caregivers About Seizures

- 1. Loosen tight clothing, especially around the neck.**




- 2. Make sure the person doesn't fall.**

● Hold the person steady if he or she is on a chair, couch, or bed.

● If the person is standing, get them to the ground safely.

- 3. Turn the person and their head to the side so that anything in their mouth, even spit, doesn't block the throat.**



- 4. Don't put anything in the person's mouth; you may get bitten.**



- 5. If you know cardiopulmonary resuscitation or CPR, check the heartbeat in the neck.**

● If there is no pulse, start CPR. Call 911.

- 6. Listen for breathing at the mouth.**

● Extend the person's neck if they are having a hard time breathing.

● If the person isn't breathing, start CPR. Seal your lips over the person's mouth and breathe two quick breaths.

● Seal your lips over the person's mouth and breathe two quick breaths.

● Continue breathing into their mouth every 5 seconds until the person starts breathing on their own.

● Call 911.




- 7. If this is the first seizure after a TBI, call the person's doctor for advice.**

- 8. IF THE SEIZURE DOES NOT STOP AFTER 3 MINUTES, CALL 911.**

- 9. If the seizure stops within 3 minutes, call the person's doctor.**

- 10. If the person doesn't return to normal within 20 minutes after the seizure, call 911.**



The content of this infocomic was adapted from the factsheet entitled Seizures After Traumatic Brain Injury, which was developed by Jeffrey Englander, MD, David X. Cifu MD, Ramon Diaz-Arrastia MD, and Alan Towne, MD and was updated by David X. Cifu MD, Ramon Diaz-Arrastia MD, and Alan Towne, MD, in collaboration with the Model Systems Knowledge Translation Center (<https://msktc.org>).

Source: The content in this infocomic is based on research and/or professional consensus. This content has been reviewed and approved by experts from the Traumatic Brain Injury Model Systems (TBIMS), funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR), as well as experts from the Polytrauma Rehabilitation Centers (PRCs), with funding from the U.S. Department of Veterans Affairs. The contents were also reviewed by American Indians with TBI and their families.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of the infographic were developed under grants from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant numbers 90DPKT0008 and 90DPKT0009). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this infographic do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the Federal Government.

Additional Note: This resource was originally designed to support American Indians with traumatic brain injury (TBI) with special cultural considerations for their community. It can be used to support individuals with TBI from other communities.

Copyright © 2025 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center



TBIMS
Traumatic Brain Injury Model System
Since 1987



**Traumatic Brain Injury
Resource Bundle**
for American Indians