


Free research-based resources from the
MSKTC to support individuals with

Burn Injury

Edition 14
March 2026



www.MSKTC.org/burn



Offered by the MSKTC to Support Individuals Living With Burn Injury

Edition 14
March 2026

Copyright © 2026 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.

The contents of this document were developed under grants from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR), grant number 90DP0082 and 90DPKT0009. NIDILRR is a Center within the Administration for Community Living (ACL), U.S. Department of Health and Human Services (HHS). The contents of this document do not necessarily represent the policy of NIDILRR, ACL, and HHS, and you should not assume endorsement by the Federal Government.

Model Systems Knowledge Translation Center

American Institutes for Research
1400 Crystal Drive, 10th Floor
Arlington, VA 22202-3231

www.MSKTC.org

MSKTC@air.org

Phone: 202-403-5600

TTY: 877-334-3499

Connect with us on



Facebook: Burn Injury Model Systems



X social media: Burn_MS



Instagram: <https://www.instagram.com/burnmodelsystem/>

About the Model Systems Knowledge Translation Center

The Model Systems Knowledge Translation Center (MSKTC) summarizes research, identifies health information needs, and develops information resources to support the Model Systems programs in meeting the needs of individuals with spinal cord injury (SCI), traumatic brain injury (TBI), and burn injury (Burn). The health information offered through the MSKTC is not meant to replace the advice from a medical professional. Users should consult their health care provider regarding specific medical concerns or treatment. The current MSKTC cycle is operated by American Institutes for Research® (AIR®). The Center is funded by the U.S. Department of Health and Human Services (HHS), National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR) under grant number 90DPKT0009. To learn more, visit <https://www.msktc.org>.

About the Burn Model Systems Program

The Burn Model System (BMS) Program is funded by NIDILRR in the U.S. Department of Health and Human Services. The BMS began in 1994 and continues its mission to improve care and outcomes for individuals with burn injuries. Currently, there are four BMS centers, each providing a multidisciplinary system of rehabilitation care, including emergency medical, acute medical, and post-acute services. In addition to providing direct services, these centers play a pivotal role in building the national capacity for high-quality treatment and research serving persons with burn injuries, their families, and the communities in which they reside. To learn more, visit <https://msktc.org/burn/model-system-centers>.

About the National Data and Statistical Center for the Burn Model Systems

The National Data and Statistical Center for the Burn Model Systems (the National BMS Data Center) advances medical rehabilitation by increasing the rigor and efficiency of scientific efforts to assess the experiences and outcomes of individuals with burn injury. Specifically, the National BMS Data Center:

- Maintains the national longitudinal database (BMS Database) for data submitted by each of the Burn Model Systems Centers (BMS Centers).
- Facilitates the entry of high-quality, reliable data in the BMS Database by providing training and technical assistance to BMS Centers.
- Facilitates the entry of high-quality data collected from database participants of all racial and ethnic backgrounds by providing knowledge, training, and technical assistance to the BMS Centers on culturally appropriate methods of longitudinal data collection and participant retention.
- Supports rigorous research conducted by BMS Centers and investigators from outside of the BMS network who are analyzing data from the BMS Database by making statistical and other methodological consultation available.
- Improves the efficiency of the BMS Database operations through collaboration with other entities, such as the National Data and Statistical Center for Traumatic Brain Injury Model Systems, the National Data and Statistical Center for Spinal Cord Injury Model Systems, the Model Systems Knowledge Translation Center (MSKTC), and the American Burn Association.
- Improves reports for the public from the BMS Database.

The National BMS Data Center is operated by the University of Washington Department of Rehabilitative Medicine and is funded by the National Institute on Disability, Independent Living, and Rehabilitation Research.

<http://burndata.washington.edu/>

Contents

Overview of Burn Injury Products Offered Through the MSKTC	4
Opportunities to Participate in MSKTC Activities	5
Collection of Burn Factsheets Offered by the MSKTC.....	5
Amputation After Burn Injury	6
Delirium After Burn Injury	11
Employment After Burn Injury	16
Exercise After Burn Injury	22
Going Back to School After a Major Burn Injury.....	25
Healthy Eating After Burn Injury—For Adults.....	30
Healthy Eating After Burn Injury—For Kids	34
Help Your Child Recover—Build Your Child’s Resilience After a Burn Injury.....	38
Itchy Skin After Burn Injury.....	43
Laser Therapy for Burn Scars	47
Managing Pain After Burn Injury	50
Outpatient Opioid Management for Adult Burn Survivors.....	55
Post-Traumatic Stress Disorder (PTSD) After Burn Injury	59
Psychological Distress After Burn Injury	63
Scar Management After Burn Injury.....	68
Sexuality and Intimacy After Burn Injury.....	72
Sleep Problems After Burn Injury	76
Social Interaction After Burn Injury	81
Sun Protection After a Burn Injury	85
Temperature Sensitivity After Burn Injury	90
Understanding a Burn Injury	94
Understanding and Improving Body Image After Burn Injury	100
Wound Care After Burn Injury	107

Overview of Burn Injury Products Offered Through the MSKTC

The Model Systems Knowledge Translation Center (MSKTC) offers a variety of free resources on burn injury to support people with disabilities and their caregivers, researchers, practitioners and clinicians, and policy makers. The following resources are available at www.MSKTC.org/Burn.

Burn Factsheets

The MSKTC collaborates with Burn Injury Model Systems to produce evidence-based and consumer-friendly factsheets.

Burn Slideshows

The MSKTC develops traditional and narrated slideshows based on information from select factsheets. These resources are developed for users who prefer a format with more images, less text, or an audio option.

Burn Hot Topic Module

The MSKTC developed a Hot Topic Module, which is a collection of resources such as videos, factsheets, and slideshows to support individuals who live with Burn Injury.

Burn Quick-Turnaround Reviews

The MSKTC provides quick reviews of Model Systems research studies funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR). Those reviews offer timely summaries of newly released Model System research studies using lay language for easy access.

Burn Systematic Reviews

The MSKTC collaborates with Model System programs to conduct systematic reviews on high-priority health topics to inform clinical practice. Results of a systematic review provide the best information for making decisions about treatment, practice, or behavior.

Burn Database

The MSKTC maintains a database of over 350 Burn citations and abstracts of studies funded by NIDILRR.

Burn Infocomics

The MSKTC translates contents from the consumer factsheets and presents them in a comic-styled storyboard format.

Burn Audio Factsheet

The MSKTC narrates the consumer factsheets to develop audio factsheets to enhance user accessibility.

Burn Infographics

Data and information presented visually for easy comprehension.

Opportunities to Participate in MSKTC Activities

Engaging people with lived experience in our work helps make the MSKTC resources more useful. Periodically, we recruit individuals with traumatic brain injury and their family members, caregivers, and clinicians to participate in MSKTC activities. We sometimes offer a small financial token of appreciation to participants. Examples of activities include:

- ◆ Providing feedback on factsheets
- ◆ Offering input on videos
- ◆ Participating in a conversation about the needs of caregivers

If you are interested in learning more about these activities or participating, please email MSKTC@air.org or call 202-403-5600.

Collection of Burn Factsheets Offered by the MSKTC

This section contains Burn factsheets offered by the MSKTC as of March 2026. These and other resources are available at no charge on [MSKTC.org/Burn](https://www.msktc.org/Burn)

- ◆ Amputation After Burn Injury
- ◆ Delirium After Burn Injury
- ◆ Employment After Burn Injury
- ◆ Exercise After Burn Injury
- ◆ Going Back to School After a Major Burn Injury
- ◆ Healthy Eating After Burn Injury—For Adults
- ◆ Healthy Eating for Kids With Burn Injury
- ◆ Help Your Child Recover—Build Your Child’s Resilience After a Burn Injury
- ◆ Itchy Skin After Burn Injury
- ◆ Laser Therapy for Burn Scars
- ◆ Managing Pain After Burn Injury
- ◆ Outpatient Opioid Management for Adult Burn Survivors
- ◆ Post-Traumatic Stress Disorder (PTSD) After Burn Injury
- ◆ Psychological Distress After Burn Injury
- ◆ Scar Management After Burn Injury
- ◆ Sexuality and Intimacy After Burn Injury
- ◆ Sleep Problems After Burn Injury
- ◆ Social Interaction After Burn Injury
- ◆ Sun Protection After Burn Injury
- ◆ Temperature Sensitivity after Burn Injury
- ◆ Understanding a Burn Injury
- ◆ Understanding and Improving Body Image After Burn Injury
- ◆ Wound Care After Burn Injury

Amputation After Burn Injury

September 2025

<https://msktc.org/burn/factsheets>

BURN Factsheet

This factsheet explains what patients can expect from amputation after a burn injury. It also notes the role that family members and the health care team play in improving the patient's quality of life after amputation.

The Model Systems Knowledge Translation Center works with Burn Model System (BMS) centers to provide free research-based rehabilitation resources for people living with burn injury. (See <https://msktc.org/burn> for more information.) This factsheet has been approved by experts from the BMS centers.

Introduction

Amputation is the whole or partial loss of a body structure (leg, arm, finger, ear, etc.). Amputation after a burn injury is rare, but it does occur in around 2% of burn survivors. Amputation may help to improve function after a severe burn injury because the patient does not need to go through many years of debridement, limb-salvaging surgeries, and pain.



Why Is Amputation Needed?

Many amputations result from a physically traumatic event, such as a burn injury. People with severe heat and electrical burns are at higher risk of amputation. Severe burns increase the chance of the following:

- Limbs, digits, or other body parts (e.g., nose, ears, eyes) are no longer functional or do not respond to surgical or therapeutic interventions.
- Sepsis, which is a severe blood infection.

An amputation can remove infection and improve healing, the chance of survival, and the chance of better outcomes in some patients. Sometimes patients and family members must make the difficult decision about whether amputation is the right choice for them. Clinicians can provide guidance and support to help them navigate this decision.

What to Expect After Amputation

Patients with a burn-related amputation can experience physical and emotional changes and challenges. Most will participate in a rehabilitation program and adjust to a new lifestyle.

Physical Outcomes

Range of Motion

Range of motion refers to how much a joint can move. Scarring can limit a patient's range of motion. Burn patients have a high risk of developing the following:

- Hypertrophic scarring (thick, deep scars in the skin)
- Contractures (thick, deep scars that cross over joints)

Engaging in range-of-motion exercises and rehabilitation therapy soon after amputation may decrease the effects of scarring. Having good range of motion helps patients with activities of daily living (getting dressed, brushing your teeth, etc.), functional activities, recovery, and potentially using a prosthesis in the future.



Feeling and Sensitivity of Skin

The injured skin around an amputation site may

- Be extra sensitive to touch (e.g., the feeling of different fabrics) or to hot and cold temperatures
- Have decreased or increased sensitivity, which can increase the chance of skin breakdown over time



Burn injury over muscles and tendons may heal slowly and require grafting or plastic surgery, which can delay prosthetic fitting and rehabilitation. Three parts of early rehabilitation will help to prepare the skin and the rest of the limb for the prosthesis:

- Using compression wraps or bandages to manage edema (swelling) and help shape the residual limb
- Stretching to maintain or increase the range of motion and reduce muscle and scar tightness
- Doing range-of-motion exercises, which are important to support functional use of the extremity with or without a prosthesis

Pain and Pain Management

Three types of pain are common among burn survivors who have sustained an amputation:

- **Residual limb pain (RLP).** RLP is pain at the amputation site and in areas of the remaining limb. RLP is common during the first few weeks after amputation. This type of pain tends to decrease as the remaining limb heals.
- **Phantom limb sensation (PLS).** With PLS, patients think they feel sensations—such as tingling or itching—in the part of the limb or digit that is no longer there.
- **Phantom limb pain (PLP).** PLP is a sharp, shooting, or burning pain. The level of pain varies, and it tends to go away over time. PLP is common after amputation and does not prohibit success with a prosthesis.

The health care team should prescribe an individualized program to manage pain from the burn injury and pain associated with an amputation. The team should monitor or adjust pain management programs during the prosthetic fitting process. These programs can include oral or topical medications (medications that go on the skin), emotional and rehabilitation therapy, mirror therapy, and prosthetic adjustment. Pain from the burn or amputation may still occur years after amputation. Patients should consult with their medical team to explore ways to manage pain symptoms. The [Managing Pain After Burn Injury](#) factsheet contains more details about managing pain.

Pruritus

Itchy skin, or “pruritus,” is a common and normal side effect of a healing burn injury. Itching may interfere with a patient’s daily activities and overall health but should decrease over time. The [Itchy Skin After Burn Injury](#) factsheet gives a detailed overview of possible treatments and therapies to help with itchy skin.

Emotional Impact

Amputation after a burn injury is a life-changing event, and those who experience amputation may go through emotional changes and distress. Many have reported becoming less social and disliking their body image. Patients may also feel sad, angry, or anxious, or experience a loss of independence. Early support from family members, friends, and the health care team should focus on protecting and improving the survivor's mental health.



Peer support has been shown to be tremendously valuable in the recovery of burn survivors who experience amputation. There are many local and national burn and amputee peer support groups that offer in-person or virtual support. The linked resources below offer more details on mental distress after a burn injury and accessing peer support:

- [Psychological Distress After Burn Injury](#)
- [Post-Traumatic Stress Disorder \(PTSD\)](#)
- [Understanding and Improving Body Image After Burn Injury](#)
- [Burn Peer Support: Phoenix Society for Burn Survivors](#)
- [Support Groups and Peer Support: Amputee Coalition](#)

Rehabilitation

Physical and/or Occupational Therapy

The goals of physical and occupational therapy include pain management and increasing the range of motion and strength of burn survivors, all with the goal of maximizing a return to independence and activities of daily living after a burn injury. Activities of daily living include tasks such as brushing your teeth, getting dressed, showering, getting in and out of bed, etc. Therapists will also help survivors learn to become independent in putting on, using, and taking off their prosthetic.

Prosthesis

The need for a prosthesis varies by type of amputation, desired use (e.g., cosmetic), and functional need. Survivors tend to be fitted for a prosthesis 1–6 months after amputation and should seek a referral from their health care team to begin the prosthetic fitting and rehabilitation process. However, there is no set timeline for a prosthetic, and patients may get fitted for one later. The [Amputee Coalition](#) offers more information about prostheses.

Exercise

Exercise—beginning as early as possible after amputation—is crucial for overall health and well-being. However, survivors may not view or participate in exercise in the same way as they did before their injury. Survivors should speak with their health care team to develop an exercise plan that fits their needs. There are many adaptations that can be made to continue activities enjoyed prior to the injury.



Life After Amputation

Many burn survivors who experience amputation lead healthy and fulfilling lives. They regain their independence and adapt to doing the activities they love, perhaps just in a different way. Remember, recovering from a burn injury can take time, and everyone progresses at their own pace. Survivors can work with their health care team to develop achievable goals and functional milestones for their recovery process. Living and thriving with a burn injury is possible!

Additional Resource

The Phoenix Society for Burn Survivors is the leading national nonprofit organization dedicated to empowering people affected by burn injury. It serves burn survivors, loved ones, burn care professionals, researchers, and anyone else committed to supporting the burn community and building a safer world. The organization provides in-person and online peer support, on-demand resources that are available 24/7, connections to a network of support organizations, and so much more. For more information, visit www.phoenix-society.org.

References

Bartley, C. N., Atwell, K., Purcell, L., Cairns, B., & Charles, A. (2019). Amputation following burn injury. *Journal of Burn Care & Research*, 40(4), 430–436.
<https://doi.org/10.1093/jbcr/irz034>



Ferguson, J. R., & Blanck, R. (2011). Prosthetic management of the burn amputation. *Physical Medicine and Rehabilitation Clinics of North America*, 22(2), 277–299, vi. <https://doi.org/10.1016/j.pmr.2011.03.001>

Kennedy, P. J., Young, W. M., Deva, A. K., & Haertsch, P. A. (2006). Burns and amputations: A 24-year experience. *Journal of Burn Care & Research*, 27(2), 183–188.
<https://doi.org/10.1097/01.bcr.0000203492.89591.a1>

Modest, J. M., Raducha, J. E., Testa, E. J., & Eberson, C. P. (2020). Management of post-amputation pain. *Rhode Island Medical Journal*, 103(4), 19–22. <http://www.rimed.org/rimedicaljournal/2020/05/2020-05-19-pain-modest.pdf>

Şimsek, N., Öztürk, G. K., & Nahya, Z. N. (2020). The mental health of individuals with post-traumatic lower limb amputation: A qualitative study. *Journal of Patient Experience*, 7(6), 1665–1670.
<https://doi.org/10.1177/2374373520932451>

Authorship

Amputation After Burn Injury was developed by Lauren Shepler, MPH; David Crandell, MD; Laura Gosselin, OT; Megan Tinney, PT, DPT; Karen Kowalske, MD; and Jeffrey Schneider, MD, in collaboration with the Model Systems Knowledge Translation Center (MSKTC).

Source: The content in this factsheet is based on research and/or professional consensus. This content has been reviewed and approved by experts from the Burn Model System (BMS) centers, funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR) and the Phoenix Society for Burn Survivors. The content of the factsheet has also been reviewed by individuals with burn injury and/or their family members.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider about specific medical concerns or treatment. The contents of this factsheet were developed under grants from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant numbers 90DPKT0009, 90DPBU0001, 90DPBU0008). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Recommended citation: Shepler, L., Crandell, D., Gosselin, L., Tinney, M., Kowalske, K., & Schneider, J. (2025). *Amputation after burn injury*. Model Systems Knowledge Translation Center (MSKTC). <https://msktc.org/burn/factsheets/amputation-after-burn-injury>

Copyright © 2025 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Delirium After Burn Injury

April 2020

www.msktc.org/burn/factsheets

BURN Factsheet

This factsheet explains what delirium is and what you or your friends and family can do about it.

What you need to know about delirium

- What is delirium?
- What are the symptoms of delirium?
- After a burn injury, who develops delirium in the hospital?
- What can the health care team do to help?
- Does delirium go away?
- What are the long-term effects of delirium?
- What can you do to help someone who has delirium?
- Frequently asked questions about delirium

What is delirium?

Delirium (dee-leer-e-um) is a 'disturbance in mental abilities that causes confused thinking and reduced awareness of one's environment.'¹ It is usually caused by an underlying health problem, such as an infection. Other causes include changes in the sleep cycle, an illness, medicines, or drug withdrawal. Delirium does not mean that your loved one is "crazy" or has a permanent mental illness.

Three types of delirium have been identified:^{1, 2}

- **Hyperactive** delirium. This is likely the easiest type to recognize. Signs include feeling restless or agitated; people may pace. People may have rapid mood changes. They may also hallucinate, or see things that are not there. They may not cooperate with care and pull at intravenous (IV) lines and tubes or try and get out of bed.
- **Hypoactive** delirium. Signs of this type may include inactivity (slow and/or limited activity). People may feel sluggish or more tired than normal. They may seem to be in a daze and may not recognize loved ones.
- **Mixed** delirium. This type includes both hyperactive and hypoactive signs and symptoms. The person may quickly switch back and forth between these two states.

What are the symptoms of delirium?^{1, 2}

Symptoms of delirium may begin over a few hours or a few days. They may come and go throughout the day. Other times, people may have no obvious symptoms. Symptoms tend to be worse at night when it is dark. Some of the main symptoms include the following.

The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <http://www.msktc.org/burn/model-system-centers> for more information).



Model Systems
Knowledge Translation
Center

Reduced awareness of the environment. This includes:

- Not being able to stay focused.
- Being easily distracted.
- Being withdrawn, with little or no activity or little response to the environment.

Poor thinking skills. This includes:

- Having poor memory and attention. A person may not recall instructions. He or she may ask for questions to be repeated.
- Being disoriented. A person may not know where he or she is or what is happening.
- Problems with speaking. This may include rambling or nonsense speech.
- Problems with reading or writing.

Changes in behavior. This may include:

- Seeing things that are not there.
- Being restless, agitated, or even aggressive.
- Calling out. This may include moaning or saying words that do not make sense.
- Being withdrawn; this is common in older adults.
- Having or showing little or no feeling or emotion.
- Not sleeping well.
- Reversing day and night cycles. This causes trouble sleeping at night and daytime drowsiness.

Emotional disturbances or changes. These include:

- Having a wide range of emotions. People may feel anxious, scared, or paranoid. They may also feel sad, cranky, or angry.
- Having sudden mood swings that cannot be predicted.
- Having personality changes.

After a burn injury, who develops delirium in the hospital?

Only a few studies have looked at delirium in burn patients. One study found that as many as 77% of burn patients who need a breathing tube develop delirium in the intensive care unit (ICU).³ Delirium may also develop after an operation. About 15% of burn patients develop delirium after an operation.⁴ Older age is a risk factor for developing delirium in the hospital. In some cases, delirium may have no known cause.

What can the health care team do to help?

Doctors and nurses use several screening tools daily to help identify delirium. They may ask the same questions when they see the patient each day, or even during every nursing shift.

The health care team knows that delirium may develop, especially in patients in the ICU. They will try to prevent it. This may include having the patient walk around the nursing unit or sit up in a chair. Patients should walk as soon as they can and often (make sure the burn team okays this type of activity). The team may also have the patient stay awake during the day and look for ways to help the patient sleep at night. The team may ask you to help with this by adjusting lights and noise in the room at certain times of day.

The team will try to manage your loved one's pain from their burn injury without over- or under-treating them.

The team may limit medicines that can make people feel confused. If your loved one is breathing with the help of a breathing machine, the doctors and nurses will try to decrease the use of medicines that make your loved one drowsy.

They will treat infections, correct fluid and electrolyte imbalances, and promote normal sleep and wake cycles.

They will try to limit interruptions by grouping nighttime care (medication administration, vital signs, and other assessments).

The care that a patient with delirium receives or the strategies used by the health care team to treat or prevent delirium may differ.

What can you do to help someone who is experiencing delirium?

The health care team may ask you about your loved one's medical history and usual behavior. Often, patients with delirium cannot provide an adequate medical history. Helping the team to understand how your family member's mental state is different from usual is a big help. It is also important that you tell your loved one's doctors and nurses if you notice a sudden change in the patient's behavior or emotions.

It is helpful to tell the health care team about your loved one's home routine and habits. This may include what time they wake up and go to sleep, their hobbies, and their likes and dislikes. Doing familiar tasks can help make a long hospital stay feel more normal.

Minimize daytime naps to help reset the day/night sleep cycle. Keep the lights on and the blinds open during the day. Turn the TV and lights off at the same time every night. Using an eye mask, ear plugs, or headphones (that play calming music) can help drown out the noise of the hospital. These habits can also help improve the patient's sleep.

Make sure to bring your loved one's glasses and hearing aids to the hospital. This will help them recognize you and their surroundings, which improves delirium. Bring a few photos of the patient and their loved ones, including pet photos, to help reorient them. Bringing in a favorite toy or stuffed animal may also help the child who experiences delirium.

Do not argue with your loved one if he or she is confused. This will lead to frustration. Instead, gently reassure the patient and remind them that they are safe in the hospital. It may also help to remind your loved one of the date, day of the week, and year. Keep instructions simple and give your loved one time to process directions. If you get frustrated, it is okay to step away from the bedside to take a break.

When your loved one is delirious, limit noise and visitors. Identify a small group of trusted friends or family members who can be with the patient to reassure them. During these quiet times, limit talking, touching, or other stimulus.

Does delirium go away?

For most patients, delirium gets better when they get treatment for the underlying cause of the change in their mental state. This could be a medical issue, a sleep disturbance, or side effects from medicine.

What are the long-term effects of delirium?

For some patients, the effects of delirium linger after their hospital stay. These effects may not be obvious. Patients may be much improved but still have problems with certain tasks that require focused thinking.

In a study of burn patients treated at an inpatient rehabilitation facility after their hospital stay, about 25% of patients had some difficulty with memory or thinking through a new problem when they left the rehabilitation hospital.⁵

Studies with non-burn patients have found that patients who had more days of delirium while in the ICU had the most severe long-term issues with their cognition⁶ and some have trouble with activities of daily living and functioning.⁷

Frequently Asked Questions About Delirium

1. **Is delirium a kind of dementia?** No, but it may be hard to tell the difference between dementia and delirium sometimes. A person can have both. 'Dementia is the loss of cognitive functioning—thinking, remembering, and reasoning—and behavioral abilities to such an extent that it interferes with a person's daily life and activities.'⁸
2. **How can I tell the difference between dementia and delirium?** Signs of delirium start over a few hours or a few days; dementia begins with minor symptoms that slowly get worse over time. With delirium, staying focused is difficult. A person in the early stages of dementia is generally alert. Symptoms of delirium may shift or change often throughout the day. With dementia, people have fairly consistent memory and thinking skills during the course of a day.

Comparison between delirium and dementia⁹

	Delirium	Dementia
Timing of onset of symptoms	Develops quickly, in hours or days	Develops over months or even years
Impact on memory or thinking	Delirium can make memory and thinking problems worse	Dementia is a disturbance of thinking
Length of illness/symptoms	Usually clears up after a few days or weeks	Usually a permanent condition

3. **Is delirium a disease?** No, it is a group of symptoms.
4. **What is usually the first sign of delirium?** Sudden confusion about time and place (where they are).
5. **Can delirium occur at any age?** Yes, delirium can occur at any age to include children, but it is more common in the elderly.

Additional Resources

Critical Illness, Brain Dysfunction, and Survivorship (CIBS) Center: <https://www.icudelirium.org>. Accessed January 21, 2020.

Patient Education: Delirium (Beyond the Basics): https://www.uptodate.com/contents/delirium-beyond-the-basics?topicRef=16994&source=see_link. Accessed January 21, 2020.

References

- ¹ Mayo Clinic. (n.d.). *Delirium: Symptoms and causes*. Retrieved from <https://www.mayoclinic.org/diseases-conditions/delirium/symptoms-causes/syc-20371386>. Accessed February 27, 2019.
- ² Gleason, O. C. (2003). Delirium. *American Family Physician*, 67(5), 1027–1034.
- ³ Agarwal, V., O'Neill, P. J., Cotton, B. A., Pun, B. T., Haney, S., Thompson, J., Kassebaum, N., Shintani, A., Guy, J., Ely, E. W., & Pandharipande, P. (2010). Prevalence and risk factors for development of delirium in burn intensive care unit patients. *Journal of Burn Care and Research*, 31(5), 706–715.
- ⁴ Guo, Z., Liu, J., Li, J., Wang, X., Guo, H., Ma, P., Su, X., & Li, P. (2017). Postoperative delirium in severely burned patients undergoing early escharotomy: Incidence, risk factors, and outcomes. *Journal of Burn Care and Research*, 38(1), e370–e376.
- ⁵ Hendricks, C. T., Camara, K., Violick Boole, K., Napoli, M. F., Goldstein, R., Ryan, C. M., & Schneider, J. C. (2017). Burn injuries and their impact on cognitive-communication skills in the inpatient rehabilitation setting. *Journal of Burn Care Research*, 38(1), e359–e369.
- ⁶ Girard, T. D., Jackson, J. C., Pandharipande, P. P., Pun, B. T., Thompson, J. L., Shintani, A. K., Gordon, S. M., Canonico, A. E., Dittus, R. S., Bernard, G. R., & Ely, E. W. (2010). Delirium as a predictor of long-term cognitive impairment in survivors of critical illness. *Critical Care Medicine*, 38(7), 1513–1520.
- ⁷ Brummel, N. E., Jackson, J. C., Pandharipande, P. P., Thompson, J. L., Shintani, A. K., Dittus, R. S., Gill, T. M., Bernard, G. R., Ely, E. W., & Girard, T. D. (2014). Delirium in the intensive care unit and subsequent long-term disability among survivors of mechanical ventilation. *Critical Care Medicine*, 42(2), 369–377.
- ⁸ National Institute on Aging. (2017). *What Is dementia? Symptoms, types, and diagnosis*. Retrieved from <https://www.nia.nih.gov/health/what-dementia-symptoms-types-and-diagnosis>. Accessed April 9, 2019.
- ⁹ Critical Illness, Brain Dysfunction, and Survivorship (CIBS) Center. (n.d.). *Patients and families overview*. Retrieved from <https://www.icudelirium.org/patients-and-families/overview>. Accessed January 21, 2020.

Authorship

Delirium after Burn Injury was developed by Gretchen J. Carrougher, MN, RN; Taylor Powell, BSN, RN (Northwest Regional Burn Model System; UW Medicine Regional Burn Center, Seattle, WA); Jeffrey C. Schneider, MD (Boston-Harvard Burn Model System, Boston, MA); and Kimberly Roaten, PhD, CRC (North Texas Burn Model System, Dallas, TX) in collaboration with the Model Systems Knowledge Translation Center.

Source: The content in this factsheet is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems (BMS), funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR).

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this factsheet were developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Copyright © 2020 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Employment after Burn Injury

March 2016

www.msktc.org/burn/factsheets

BURN Factsheet

This factsheet covers returning to work after burn injury, including how to know when you're ready to return to work and accessing workplace accommodations.

Returning to work after a burn injury can be an important phase of recovery that helps you return to a routine. Work not only provides you with an income and other benefits, but can also give you a sense of purpose and confidence that is critical in maintaining a higher quality of life. Returning to work, even if you were not injured on the job, can be accompanied by a confusing range of emotions, such as feeling both excitement and anxiety.

- You may not feel you are physically or mentally ready to return to work and do not have the stamina to work the same amount of hours or perform the same physically demanding tasks that you did before you were injured.
- If you were injured on the job, then returning to the same place of injury and/or the same tasks that you were doing when you were injured, may lead to fears or anxiety related to the injury.

Many people have found it was helpful for them to start work gradually, even before they were fully recovered. This is because returning to work can be emotionally and physically therapeutic.

There are many resources available to help you return to work. Some of these resources vary according to your state law and whether or not you were injured on the job. Your eligibility for resources depends on your unique circumstances such as the severity of your burn injury and the demands of your job.


The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <http://www.msktc.org/burn/model-system-centers> for more information).

Remember that you are not alone. It is important to get assistance from the burn center, your employer, or other community resources to help you take the steps necessary to return to work safely and successfully.

Are You Ready to Return to Work?

There are some important things that you can do to help make the process of returning to work easier:

- See your health care providers regularly and follow through with treatment. Success with getting back to work requires active participation on your part.
- Talk with your health care providers about your readiness to return to work. He or she can help assess your current strengths and limitations and a reasonable time frame for returning to work.
- Spend time during your recovery period to focus on what you need to do emotionally and physically to return to work. This might include doing tasks around the house to build your strength and stamina, participating in an exercise program, or wearing work boots around the house to get comfortable in them again, particularly if you had a burn to the foot or leg.
- Meet with someone who has returned to work after an injury to get support and guidance.



Model Systems
Knowledge Translation
Center

- Build a supportive network to help you role-play such things as interviews and educating employers and co-workers about your situation.
- Getting prepared may also include counseling to help you cope with psychological and emotional issues you have about returning to work.
 - You may feel uncomfortable about your change in appearance due to your burn injury, especially if your burns are highly visible such as on your face or hands.
 - You may want to learn how to handle questions from your coworkers or employer about your injury and how it happened, or about the way the burn has changed the way that parts of your body look (e.g., scars, pigmentation).
 - If you were injured on the job, treatment for Post-traumatic Stress Disorder (PTSD) symptoms will be important. Treatment might focus on being prepared to return to the place and activities where you were working and other reminders of your accident, such as coworkers who were there when you were injured. It will also include managing nightmares, flashbacks and other symptoms of PTSD.
 - Some people have found it helpful to get in touch with some coworkers they are looking forward to working with again.
- Stay in touch with your employer and express your interest in getting back to work. Keep your employer informed about your recovery to ensure your job will be waiting for you when you are ready to return.
- Ask your health care provider to clearly evaluate your current work restrictions and a time frame for when you may be able to return to work.
- Ask him or her to write a note to your employer with a timeframe and list of supports (workplace accommodations) you may need so that your employer can plan accordingly.
- If you are not able to return to your regular job for a while, talk to your employer about jobs that you may be able to do during your recovery.

Work Conditioning Programs

If your medical team determines that you are ready to return to work but you and they feel you lack the stamina needed to do your work, you may participate in a return to work program that will gradually build your stamina for work, or be encouraged to do fitness training. If your injury happened at work, you might also be referred to participate in a work conditioning (work hardening) program. These are specialized programs run by occupational and physical therapists that aim to improve your strength and endurance to perform your job tasks. Discuss options for improving your endurance for work with your health care providers since there are advantages and disadvantages to various approaches.

Workplace Changes to Help You Return to Work

Your medical team may ask your employer for a job description when you are ready to return to work. This job description is usually provided by the human resources department at your place of employment. The job description helps your medical team determine if you are physically capable of meeting the demands of your job, or if you need changes made at your workplace to accommodate your disability (accommodations). It is OK to ask your employer for accommodations to help you return to work. The Department of Human Resources at your place of work is a good place to start this dialogue. Whether you were burned on the job or not, employers are legally obligated under the Americans with Disabilities Act and many state laws, to provide you with reasonable accommodations as long as it will not cause the business any undue hardship.

- Under the law, it is your responsibility to make the request for accommodations.
- Be prepared to advocate for yourself or find someone to advocate for you because your employer may not be familiar with the laws that require job accommodations.
- Your medical team may provide you with guidance and supporting paperwork to help you receive job accommodations.
- You have the right to refuse a job assignment that violates your doctor's stated restrictions.

Typical Workplace Accommodations Include:

- A modified work schedule (4 hours per day in the beginning, with a gradual increase to full time).
- Working in a clean and dry environment.
- Limits on how much you can lift.
- Limited exposure to extreme temperatures (avoiding extreme heat or prolonged periods of cold).
- Altering work schedules to accommodate outpatient physical therapy and other medical appointments.
- Special software and hardware so a person with limited or no hand function can work on a computer.
- Adaptive equipment to help you perform your job tasks.
- Consider having your first day at work be in the middle of the week, rather than on a Monday. This short week will be less physically demanding for you.

These accommodations can make the initial transition back to work less demanding on you, and over the long haul, make it possible for you to succeed in your job. You may also need more recovery time when you get home at the end of the workday. It may be helpful to educate your employer and co-workers about burn injuries so they are more understanding to your needs and have realistic expectations of your performance.

On-The-Job-Injury

If you were injured at work, you are likely covered by your state's workers' compensation insurance. Your case manager is responsible for making sure you get the benefits you are entitled to because of your on-the-job injury. Your case manager can help you get the treatment you need and make the transition back to work go more smoothly. He or she can also be a great resource in helping you to navigate the complex workers' compensation system. It is also important that you know your rights and be prepared to advocate for yourself. State workers' compensation insurance differs by state. Your case manager may not provide you with all the information about what workers' compensation benefits you are entitled to such as mileage compensation for doctor appointments.

Here are some things you can do to make this process run smoothly:

- Be sure to keep all your paperwork.
- It is important for you to know your claim number and to be in touch with your case manager regarding your treatment and your plan to return to work.
- Always keep your medical appointments to ensure your claim remains open.
- Remind your health care provider to send in the required workers' compensation paperwork so your claim is not closed before you are ready to return to work.



Model Systems
Knowledge Translation
Center

- Do not ignore calls, mail or other forms of communication from your workers' compensation program. Some of these communications may require you to take action.
- Make sure your case manager knows the name and address of your health care provider. Good communication between your health care provider, you, and your claims manager is essential to the smooth delivery of benefits.

Independent Medical Evaluation

The workers' compensation case manager or your health care provider may request an Independent Medical Evaluation (IME). These are often done when the health care provider feels that you have recovered as much as you are likely to recover. If you receive an IME, you would be referred to an outside health care provider who would independently assess if your condition is stable and no further treatment is appropriate. In that case, your claim is ready for closure.

Permanent Partial Disability Award

If your injury caused permanent loss of physical or cognitive function or a chronic psychiatric illness you will receive something called a "permanent partial disability award." Any permanent partial disability award you receive is based on the degree of physical or mental impairment on activities of daily living, not on whether you can work.

Vocational Rehabilitation

Whether you were injured on the job or not, you may need vocational rehabilitation to help you return to work.

How Does Vocational Rehabilitation Work?

If you are treated at an inpatient rehabilitation setting, vocational rehabilitation usually starts the first day you are admitted and continues after discharge, as the rehabilitation team and counselor continue to follow your employment status and provide help as necessary. Vocational rehabilitation can also take place at any time an individual with a disability needs help finding employment. If your burn injury makes it difficult for you to get or keep a job, you may try to seek assistance from a vocational rehabilitation counselor.

Your vocational rehabilitation counselor may provide the following support to help you find a job:

- Assessing your skills, limitations, health needs, work and education histories, interests and even personality style.
- Helping you develop a plan that includes specific employment-related goals and how to achieve them.
- Setting up a "trial" work situation for you with a potential employer for a specified period of time to see if you are able to do the job, if you like the job, and what accommodations you might need.
- Helping you determine whether a potential job would be a good match for you. He or she can conduct a job analysis to determine the actual kinds of tasks done in that particular job, the cognitive (mental or thinking), social and physical demands of the job, and the need for accommodations.

How You Can Find a Vocational Rehabilitation Counselor

- Ask your health care provider or burn center for a referral to a vocational counselor.
- The State Department of Vocational Rehabilitation agency (DVR) can be found in almost every state (see Resources). The DVR may also be able to help you train for a new career if you are unable to return to the job you had before your injury.

Vocational rehabilitation services vary by state. State DVRs are permitted to enter what is called an "order of selection" during difficult budget times, which means people with more severe disabilities will be given priority to receive services.

Long term disability

Due to the severity of the injury, some people may not be able to return to any type of employment. A person is eligible to receive disability payments from the Social Security Administration (SSA) if he or she is unable to perform any kind of work at a level of “substantial and gainful” activity and the disability is expected to last at least a year. The SSA pays disability benefits under two programs:

- Social Security Disability Insurance (SSDI) is a long-term disability insurance program funded by payroll taxes. In addition to disability, eligibility is based on your past work history and income. After a waiting period, SSDI recipients are eligible for Medicare.
- Supplementary Security Income (SSI) is a federal welfare program for people with disabilities who are unable to work and are poor. SSI recipients are eligible for Medicaid after one month.

If it seems unlikely that you will return to work within a year, we recommended that you apply for SSA benefits as soon as possible. If you have purchased a long term disability policy, you may also be eligible for benefits from that policy provider.

What Will Happen to Your Long Term Disability Benefits if You Return to Work?


Sometimes people are reluctant to start working because they don't want to lose their medical benefits under SSDI or SSI, or because of the extra costs of getting to or staying at work. Several federal programs have been created to help avoid this situation. Federal work incentive programs under SSDI or SSI allow people with disabilities to receive benefits and federal health care (Medicare and/or Medicaid) while still keeping some of their earnings from employment. Two federal work incentive programs that help people with disabilities secure a variety of supports such as job coaches, transportation, equipment, and work-site accommodations, include:

- PASS (Plan to Achieve Self Support) – This program (3 year max) allows funds used in achieving employment goals such as paying tuition for training courses to be excluded in calculating your monthly SSI benefits. Contact your local SSA office or for more information, go to <http://www.socialsecurity.gov/pubs/11017.html>
- IRWE (Impairment-Related Work Expenses) – IRWE benefits can be claimed for your entire working life. In this program, work-related expenses such as adaptive equipment or personal care costs can be deducted from your income so you can retain more of your SSI benefits.

The Law Protects You

The federal Americans with Disabilities Act (ADA) prohibits employers from discriminating against qualified individuals with disabilities who are able to perform the essential functions of the job with or without accommodations. To be protected under this law, you must have a disability that limits major life activities.

- An employer must make “reasonable accommodation” to your disability if it would not impose an undue hardship on the business.
- Whether an accommodation is considered a hardship depends on the business size, financial resources, nature of operation and other factors.
- When you apply for a job, an employer cannot ask you about the existence, nature or severity of your disability, but can ask you about your ability to perform certain job functions, although some elements of your disability may be visible to others.
- An employer can require you to pass a medical examination only after a conditional offer of employment are made and if it is job-related and required of all employees in similar jobs.



Model Systems
Knowledge Translation
Center

If you feel an employer has discriminated against you, contact the U.S. Equal Employment Opportunity Commission (EEOC). Call 1-800-669-4000 to find the office nearest you, or go to www.eeoc.gov. Many states have powerful disability rights laws as well.

Resources

Benefits for People with Disabilities, Social Security Administration. 1-800-772-1213. <https://www.ssa.gov/disability/>
Find your local Vocational Rehabilitation Agency and other federal and state resources, contact the Job Accommodation Network, U.S. Department of Labor. 1-800-526-7234. <http://askjan.org/cgi-win/TypeQuery.exe?902>

Equal Opportunity Employment Commission (ADA) 1-800-669-3362. <https://www.eeoc.gov/>

The Phoenix Society for Burn Survivors. 1-800-888-2876. <https://www.phoenix-society.org/>

References

Esselman, PC, Askay SW, Carrougner GJ, Lezotte DC, Holavanahalli RK, Magyar-Russell G, Fauerbach JA, Engrav LH. (2007). Barriers to return to work after burn injuries. *Archives of Physical Medicine and Rehabilitation*, Dec:88(12sup). S50-56.

Brych SB, Engrav LH, Rivara FP, Ptacek JT, Lezotte DC, Esselman PC, Kowalske KJ, Gibran NS. (2001). Time off work and return to work rates after burns: systematic review of the literature and a large two-center series. *Journal of Burn Care Research*, 22(6):401-405.

Mason, S. T., Esselman, P., Fraser, R., Schomer, K., Truitt, A., & Johnson, K. (2012). Return to work after burn injury: a systematic review. *Journal of Burn Care & Research*, 33(1):101-109.

Authorship

Employment after Burn Injury was developed by Shelley A. Wiechman, Ph.D., and Sabina Brych, B.S., in collaboration with the Model Systems Knowledge Translation Center.

Factsheet Update

Employment after Burn Injury was reviewed and updated by Lynne Friedlander, MED, Spaulding Rehabilitation Hospital in collaboration with Oswald Mondejar, Sr. VP Spaulding Rehabilitation Network; Joan Phillips, Assistant Commissioner of Vocational Rehabilitation, Commonwealth of Mass; Peter Esselman, MD, MPT, University of Washington, and James Fauerbach, PhD, Johns Hopkins. The review and update was supported by the American Institutes for Research Model Systems Knowledge Translation Center.

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. This publication was produced by the Burn Model Systems in collaboration with the University of Washington Model Systems Knowledge Translation Center with funding from the National Institute on Disability and Rehabilitation Research in the U.S. Department of Education, grant no. H133A060070. It was updated under the American Institutes for Research Model Systems Knowledge Translation Center, with funding from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this fact sheet do not necessarily represent the policy of the U.S. Department of Education or the U.S. Department of Health and Human Services, and you should not assume endorsement by the federal government.

Copyright © 2016 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Exercise After Burn Injury

February 2021

www.msktc.org/burn/factsheets

BURN Factsheet

This fact sheet explains the importance of exercise or movement after a burn injury. The information describes activities you can do to make your muscles and bones stronger and keep your joints moving.

How does a burn injury affect your body?

A burn injury causes stress to your body. Your heart and lungs may not work as well as before. Your muscles and bones may not be as strong. Remember that muscles get weak or smaller when they are not used—being on bed rest probably caused you to lose some muscle. For each day of bed rest people can lose 1% of their muscle mass.

Also, as your burns heal you may notice that your skin feels tighter. You may not be able to move your joints as far and as freely as before. This tightness and lack of movement may make it harder to take care of your everyday activities like bathing, dressing, and eating.

Why is exercise important?

The sooner you begin everyday activity, the better. Sitting up, getting out of bed, and walking will help you get out of the hospital sooner. Being active or exercising may:

- Help your breathing
- Improve your heart function
- Help your body to fight infections, like pneumonia
- Improve your flexibility and ability to move
- Lower your risk of developing scars or contractures that limit your ability to move
- Make it easier to take care of your everyday activities
- Give you a sense of well-being




What can I do?

The chart below shows the types of exercises that can benefit you. Please consult your physician before engaging in these exercises.

Type of Exercise or Activities	
Stretching	Stretching is an important part of your exercise program. • Stretching increases flexibility, which is important for preventing and treating skin contractures. • The goal of stretching is to move the joint to the point where the skin stretches. • Hold the stretch for 20 seconds to 2 minutes. Relax and repeat three times.
Aerobic activities make your heart beat faster and can make your heart, lungs, and blood vessels stronger and more fit.	Walking is an easy way to get aerobic exercise. • Walk outside or on a treadmill inside. • Start slow. • Increase the time you walk by about 1 minute per day. • Build up to walking 30 minutes to 1 hour three times a week. • You should feel as if you are working, but you should not be so short of breath that you can't talk. When cleared by your doctor, try using a stationary bike, elliptical, rowing machine or swimming.

The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <http://www.msktc.org/burn/model-system-centers> for more information).



Model Systems
Knowledge Translation
Center

Type of Exercise or Activities	
Strengthening activities make your muscles do more work than usual and make your muscles stronger.	Resistance training or muscle strengthening is exercise that uses weights, elastic bands, or your own body weight. Weights don't need to be heavy. They just need enough tension to raise your heart rate and tire your muscles. Yoga, Tai Chi, or Pilates are also ways to make your muscles stronger and keep you moving.
Recreational activities	Playing sports, gardening, and dancing are good recreational activities that can help you build strength and endurance. Children will benefit from playing games that require movement or simulated activities using video gaming technology.

Special considerations

When exercising after a burn injury, keep in mind:

- **Pain**—Talk with your doctor about the use of non-opioid medications; if fitted with custom-fit pressure garments, wear them during exercise to minimize discomfort.
- **Dry skin**—Apply creams to moisturize skin before stretching. Creams can prevent cracking or tearing of skin. Ask a family member or loved one to massage the area of tightness. Massaging the area gently before exercising can help you stretch.
- **Water**—Drink water or fluids so you do not get dehydrated during exercise.
- **Exercising in the heat**—Many people with burns are uncomfortable in the heat. Be sure to protect yourself from the sun when exercising outdoors. Cover up with a brimmed hat and long sleeves. Use waterproof sunscreen (for more information, see the MSKTC's factsheet Sun Protection After Burn Injury at <https://msktc.org/burn/factsheets/sun-protection-after-burn-injury>). Start slow and build up to longer times in the heat. Research shows that people with burns can build up a tolerance to the heat if they slowly increase exposure. Listen to your body and how you are feeling!
- **Open wounds or exposed tendons**—Talk with your doctor or therapist about what you can do to exercise safely if you have open wounds or exposed tendons.



Stretching exercises to help with tightness

The chart below shows exercises for different parts of the body where skin is tight because of a burn injury. Talk to your primary care doctor or the burn care team about the exercises that are right for you.

Face	• Look into a mirror and make facial expressions like smiling or looking surprised. • Close eyes tightly and massage skin around eyes. • Stretch your mouth open and massage the edges of your mouth. • Say the alphabet, exaggerating the letters with your mouth.
Neck	• Combine stretching your neck with face stretching. • Stretch in the opposite direction of tightness. • Lie on your back on the bed. Look up to stretch the front of your neck. As you get better, let your head jut out over the edge of the bed.
Chest	• Lie on your back with a ball or cushion in the middle of your back. – Start with your hands on your hips. – Arch your back. – Stretch both arms out to the side or over your head to increase the amount of stretch on your chest.
Shoulders	• Hold a stretch band with each hand. Use one arm to hold the other arm at the point of pull. Repeat to stretch the other shoulder. • Prop your arm on the back of the couch or chair when sitting.
Elbows	• Sit with your elbows all the way straight and your palms facing forward or up.

Hands	• Stretch each finger at the knuckle to help get the hand into a fist (see photo to the right). • For a longer stretch, wrap your hand in a fist position. • To get your hand into an open position, press down against a firm surface. – Increase the amount of stretch by using the other hand to press down on the back of the open hand.	
Knees	• To help get the knees straight, sit with your legs propped up. • Increase the amount of stretch by pressing on your thighs or knees with your hands.	
Ankles	• Standing helps stretch your ankles to get your feet flatter on the ground. • Stand on a step as if you are about to go up the steps. Lower your heel off the step.	
Toes	• Toes tend to curl up. First, massage the scar. Then use your hand to stretch the toes.	

Other Resources

The UW Medicine Regional Burn Center (Seattle, WA) has developed several short videos that demonstrate stretching exercises for specific areas of the body. Visit this YouTube site for 9 videos listed in the Burns300 series. The videos are available in both English and Spanish:
https://www.youtube.com/playlist?list=PLFEMTlzmLeUC-tONmpxadXa_7rusm_B6.



Keeping your body fit and healthy may mean going to a gym, a swimming pool, or being out in public. It's natural to be worried about how you look or how people may react to your burn scars. Check out the links below for resources that may help you.

Body Image After Burn Injury (<http://www.msktc.org/burn/factsheets/Understanding-And-Improving-Body-Image-After-Burn-Injury>)

Social Interaction After Burn Injury (www.msktc.org/burn/factsheets/social-Interaction-After-Burn-Injury)

References

Schlender ZJ, Ganio MS, Person J, et al. Heat acclimation improves heat exercise tolerance and heat dissipation in individuals with extensive skin grafts. *J Appl Physiol* 2015;119:69–76

Authorship

Exercise After Burn Injury was originally developed by Karen Kowalske, M.D., Radha Holavanahalli, Ph.D., Gretchen Carrougner, MN, RN, Oscar Suman, Ph.D., and Cindy Dolezal, P.T., MLS in collaboration with the Model Systems Knowledge Translation Center. The factsheet was reviewed by Oscar Suman, Ph.D. and Gretchen Carrougner, MN, RN in collaboration with the Model Systems Knowledge Translation Center

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this fact sheet were developed under a grant from the Department of Education, NIDRR grant number H133A110004. It was updated under the American Institutes for Research Model Systems Knowledge Translation Center, with funding from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Copyright © 2021 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Going Back to School After a Major Burn Injury

October 2020

www.msktc.org/burn/factsheets

BURN Fact Sheet

This fact sheet helps you plan for your child's return to school after a burn injury. It gives tips on when to start planning and how to prepare your child. It also offers resources and programs that may help you and your family with the changes.

Introduction

Going back to school is a very important step in a child's healing after a burn injury. Learning and being with friends is important to your child's progress. Going back to school helps your child return to a good routine and continue learning important social and academic skills. It is normal for you or your child to feel stress and be worried about going back to school. This stress can be due to changes in the way the child looks or changes in what he or she is able to do. The stress could simply be due to the time away from normal schedules. Although this step might be stressful, we encourage the child to return to his or her familiar school environment as soon as possible. For example, if your child is in a regular public school, we recommend that he or she go back to that school. Extra help and other accommodations are available in schools.



When Should My Child Go Back To School?

- After spending many days, weeks, or even months in the hospital, the sooner your child gets back to his or her routine the better.
- While each child's injury is unique, on average most children go back to school between 7 and 10 days after going home from the hospital.
- Many things affect how your child feels about going back to school – changes to the body, changes to abilities, the size of the burn, and level of self-confidence. None of these things should keep your child from going back to school as soon as possible.
- If going back to school full-time doesn't seem possible at first, starting with a schedule of half-days or every other day may be a good way to start.

The Burn Model Systems Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <http://www.msktc.org/burn/model-system-centers> for more information).



Model Systems
Knowledge Translation
Center

How and When Should I Start Planning for My Child To Return to School?

The plan for your child to go back to school should include your child and family, the school, and all members of the health care team, including the physicians, nurses, therapists, and child life specialist. Start talking about a back-to-school plan as soon as your child is medically stable and out of immediate danger. It may seem early, but the sooner the better! Starting early gives you time to put a good plan in place.

- Call your child's school and let them know that your child is in the hospital. Try to call within 2 or 3 days after the injury. If you can't make the call, ask a friend or someone on your child's health care team to call the school. Make sure the school's principal, guidance counselor or social worker, and your child's teachers are all aware of the situation. Ask them to designate one person with whom you should continue to communicate about your child's progress.
- Many children appreciate staying connected to their classmates while they are away from school. Consider asking the school to coordinate sending cards, letters, or pictures.
- Talk to your child about going back to school. Start the conversation early, while your child is still in the hospital. It will comfort your child to know that he or she will be able to get back to familiar routines and be with friends.
- Talk with the health care team about the kind of help your child will need at school and how your child's current school can best provide that help.
- Get your child's schoolwork. Ask to have schoolwork sent to the hospital. This may seem early, but having schoolwork lets your child know that he or she will get his or her life back again. It also keeps your child in contact with the outside world and gives him or her hope.



What Can I Do To Prepare My Child?

The first priority is to get your child back to doing everyday activities. Friends are also very important.

- Check with your child's health care team to see if and when you can set up short visits with close friends. If visiting in person is not possible, try other ways to get them together. Think about video chats, instant messaging, or telephone calls.
- Talk with your child about what he or she would like to share with others.
- If your child is able, suggest that he or she keep a diary or draw pictures of life in the hospital. This may help later in explaining to friends and classmates what happened.
- Children with visible wounds or scars may have questions or worries about seeing their friends for the first time. The health care team can help you prepare your child for these social interactions.
- Be supportive if your child wishes to share feelings, worries, and wishes about going back to school.
- Many burn care centers have on-site school and fun activities for children. Help your child to take part in these activities.
- For assistance with preparing you and your child for his or her return to school, consider consulting a child life specialist, psychologist, social worker, or a national organization such as the Phoenix Society for Burn Survivors.



What Are Accommodations?

Schools are required by law to provide “reasonable accommodations” for children who have been injured. The school guidance counselor or social worker is typically the person responsible for coordinating this process. These accommodations may be available to your child as he or she recovers from the injury:



- **A flexible schedule.** For example, your child may go to school for half a day and then slowly build to a full day.
- **Help taking notes.** This may include the use of a special computer or equipment or having another person take notes if their dominant hand has been affected by the injury.
- **Personal assistance.** Your child may get help with going to the bathroom, putting on a coat, or other everyday activities at school.
- **More time to take tests.** Your child may be allowed more time to take tests if he or she has trouble remembering, learning new things, or concentrating.
- **Changes to physical education.** Your child's physical education class may be adjusted or changed.
- **Help with care of the injury during school hours.** Your child may get help with stretching or putting on splints or pressure garments.
- **Visits to the school nurse.** The school nurse may be available to give your child medicine as required by you and your doctor. Many of these accommodations will be short term. Your child will need less help as he or she gets stronger and moves more easily. The steps to set up accommodations may differ from school to school. Some schools will ask you to fill out forms, while others may not. If you need an advocate, call the Special Education Department in your school district and talk to them about your child's rights and what accommodations you are requesting.

Structured vs. Informal School Re-Entry Programs

The purpose of a school re-entry plan is to provide the child, the family, and the school community with the information and support necessary to ensure a smooth transition back to school. Based on conversations with the burn care team treating your child and your child's classroom teacher(s), you should decide whether your child needs a more structured school re-entry program or simply an informal plan. A **structured program** would have a curriculum that might include sample letters to send to teachers and classmates, videos, worksheets, role plays, guest speakers and other classroom activities to raise awareness about your child's burn injury and recovery. It could be a program developed by your child's burn center or a national program such as The Journey Back developed by the Phoenix Society for Burn Survivors. You can use some or all of the components of the program as appropriate.



In contrast, an **informal school re-entry plan** may simply involve having your child's physician or another member of the burn care team, such as the child life specialist, make phone calls, send a letter, or visit the school. The hospital therapy team can help the school understand what your child can and can't do. This information can help to make sure that your child is not kept from doing school activities with the other students, such as physical education class and eating lunch. It can also make sure your child does not have to go through unnecessary infection control practices, such as not being allowed to play on the playground or not being exposed to other children in a classroom. Some things to consider when deciding whether or not you want to use a more structured program are the specific nature of your child's burn injury, whether he or she is shy and quiet, or more outgoing and comfortable with attention. It will also depend upon whether or not your child will be returning to school with pressure garments or splints, their level of endurance, and whether or not they are returning to the same school or a new school.



A good example of a back to school program is ***The Journey Back: Navigating School Reentry*** developed by the Phoenix Society for Burn Survivors:

<https://www.phoenix-society.org/resources/the-journey-back-navigating-school-reentry>

Complementary Programs

Many schools have developed other programs that may help as children return to school. Teachers and students can use the skills they learned in these programs to help your child get back into school. These programs may be called anti-bullying, disability awareness, and social and emotional learning. High schools may offer tutoring, counseling, violence prevention, and peer mentorship programs.

What Can I Do After My Child Goes Back to School?

No matter how much planning you do, you and your child may be worried about going back to school. This is natural. Reach out for help.

- **Remember to take care of yourself.** Going back to normal activities yourself is an important step for your child. It may be stressful for both of you. Make sure that you look for support for your own feelings. Keeping yourself healthy will be a big help to your child.
- **Speak up for your child.** Schools have many children who receive special services.
- **Be your child's advocate,** and ask what services are available for you or your child.
- **Check in.** In the first few weeks after your child returns to school, check in often with teachers and counselors to see how things are going. Also check in when things change, like when your child switches grades or schools.
- **Help your child work through problems.** Ask your child to be specific about problems. Then work with him or her to come up with solutions for dealing with the problems. For example, if he or she gets teased about looks, come up with different ways of handling it. Take the time to see what your child feels most comfortable doing. Then help your child practice what to say or do.



Summary

Children with burn injuries want to get back to their everyday life. Getting back to school is an important part of getting back to normal. Work with your child's health team, hospital or burn center, and school to set up a school re-entry plan so that everyone knows what your child needs. A good re-entry plan will give your child the medical and school services that he or she needs, and it will help your family adjust and heal.

References

- O'Brien, K., & Wit, S. (1985). A return-to-school program for the burned child. *Journal of Burn Care & Research*, 6(2), 108–111.
- Raquel Pan, Bruna Domingos dos Santos, Lucila Castanheira Nascimento, Lídia Aparecida Rossi, Rinie Geenen, Nancy E. Van Loey. School reintegration of pediatric burn survivors: An integrative literature review. *Burns*, Volume 44, Issue 3, 2018. Pages 494-511. <https://doi.org/10.1016/j.burns.2017.05.005>.
- Rosenstein, D. W. (1987). A school reentry program for burned children part I: Development and implementation of a school reentry program. *Journal of Burn Care & Research*, 8(4), 319–322.
- Blakeney, P., Moore, P., Meyer III, W., Bishop, B., Murphy, L., Robson, M., & Herndon, D. (1995). Efficacy of school reentry programs. *Journal of Burn Care & Research*, 16(4), 469-472.

Authorship

Going Back to School After Burn Injury was developed by Lynne Friedlander, M.Ed., and Shelley Wiechman, Ph.D., A.B.P.P., in collaboration with the Model Systems Knowledge Translation Center.

Source: The health information content in this fact sheet is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider about specific medical concerns or treatment. The contents of this fact sheet were developed under a grant (number H133A110004) from the U.S. Department of Education, National Institute on Disability and Rehabilitation Research. It was updated under the American Institutes for Research Model Systems Knowledge Translation Center, with funding from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Copyright © 2020 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Healthy Eating After Burn Injury— for Adults

November 2023

<https://msktc.org/burn/factsheets>

BURN Factsheet

This factsheet explains the importance of proper nutrition as you recover and heal from a burn injury. It describes the nutritional needs during your recovery in the hospital and at home and offers tips for balancing your diet as you heal.

This factsheet is intended to inform people with burn injury and their families about nutrition during hospitalization and after they return home. Burn injury dramatically increases your nutrition needs. The larger the burn size, the more nutrients you need to heal. A diet high in calories and protein

- Supports the immune system to decrease risk of infection
- Helps wounds heal faster;
- Maintains muscle mass
- Minimizes weight loss to support rehabilitation



How Are Nutrition Needs Determined?

A dietitian and the medical team decide how much nutrition (e.g., calories and protein) you need. They set your nutrition needs based on many factors, including your weight, height, and age, and the burn size. Vitamins and minerals, such as vitamin C, zinc, and vitamin A, are also important for healing and preventing infection. Eating a healthy, well-balanced diet will help you receive enough nutrients to support wound healing. Your medical team will let you know if you need to take any extra supplements.

How Do I Meet My Nutrition Needs While Hospitalized?

Healing from a burn injury requires more calories and protein than any other type of injury. In the hospital, a dietitian will work with you to make sure you are getting enough nutrients to heal.

After a burn injury, it is important to eat a balanced diet that includes a lot of protein in addition to other foods such as fruits, vegetables, and grains. You should eat high-protein foods at every meal and as snacks.



The Model Systems Knowledge Translation Center works with Burn Model System (BMS) centers to provide free research-based rehabilitation resources for people living with burn injury (See <https://msktc.org/burn> for more information). This factsheet has been approved by experts from the BMS centers.

Examples of high-protein foods include:

- Lean meat
- Eggs
- Dairy Products (milk, yogurt, cheese)
- Tofu
- Lentils
- Beans
- Nuts



Drinking milkshakes or smoothies may help you to meet your calorie and protein needs. If needed, the medical team may also recommend certain vitamin or mineral supplements.

You may need more nutrients than what you get from eating only by mouth. If this is the case, tube feeding can deliver more nutrients. A soft, flexible tube inserted through the nose reaches the stomach and delivers liquid formula that contains all of the nutrients needed for healing. Tube feeding will continue as long as necessary.

What if I Have Diabetes or High Blood Sugar?

After a burn injury, stress forces the level of sugar in your blood to increase. High blood sugar interferes with healing. Even if you don't have diabetes, you may have high blood sugar. Your health care team may prescribe insulin, which lowers blood sugar. Until your blood sugar improves, you might have to limit the amount of foods you consume that are high in carbohydrates, such as juice, high-sugar drinks, and desserts.

What Should I Eat at Home?

After leaving the hospital, eating a balanced diet is the best way to make sure you are staying nourished. Remember, your body now requires fewer calories than when you were hospitalized. If your burn wounds are still open, your diet should include extra protein. As you continue to heal, your nutrition needs will be like they were before the injury. At the hospital, you likely ate large meals, drank nutrition supplements, and ate a lot of snacks, so your appetite may be big when you get home. Now it is important to focus on a balanced diet. Avoid foods with little nutritional value, such as sugary beverages, desserts, candy, fatty meats, and white breads or crackers. Eat more lean meats, whole grains, vegetables, fruits, and low-fat dairy.



Ask your health care provider about exercise to help maintain a healthy weight. Exercise is important for long-term health and avoiding chronic illnesses, such as high blood pressure, diabetes, and heart disease. For more information, refer to the Exercise After Burn Injury Factsheet (<https://msktc.org/burn/factsheets/exercise-after-burn-injury>).

Tips for a Well-Balanced Diet at Home

- Eat small, frequent meals and snacks to keep from feeling so hungry that you eat too much at one time.
- Include protein with each meal and snack. Protein helps you stay full and your body stay strong. Good sources of protein are beef, chicken, pork, eggs, beans, nuts, milk, yogurt, and cheese.
- Slice fruits and vegetables for an easy snack on the go. They provide many key nutrients for healing and overall health.
- Change the flavors of the foods you eat to keep them tasty and fun. Cook with spices and herbs, such as rosemary, mint, garlic, cayenne pepper, and basil.
- Drink water during the day to stay hydrated and avoid unnecessary calories. Many drinks have fat and sugar that your body doesn't need. Read the nutrition facts on food labels.
- Choose whole-grain breads and other foods that are high in fiber. These will help keep you feeling full between meals and can support regular bowel movements.



- Try to eat only when you're hungry. Avoid eating for other reasons, such as boredom, lack of control, anger, or hopelessness. Keep a food log and track your mood to see if your eating is connected to your emotions.
- Talk to your doctor about any vitamins and supplements you would like to take at home.

Sample Menu

This menu provides about 1,750 calories and 103 grams of protein per day. This sample menu should be balanced with physical activity.

Breakfast	2 scrambled eggs 1 slice whole-wheat toast 1 tsp. butter 1 medium banana 8 oz. nonfat milk
Lunch	Turkey sandwich: 2 slices whole-wheat bread, 2 slices deli turkey, lettuce, tomato, and mustard 1 cup nonfat yogurt 8 oz. water

Snack	1 medium apple 2 Tbsp. peanut butter 8 oz. water
Dinner	3–4 oz. baked salmon 1 cup mixed vegetables ½ cup brown rice 8 oz. nonfat milk
Snack	½ cup ice cream ½ cup sliced strawberries

Additional Resources

Contact your local burn center and ask for an appointment with a dietitian who will create a nutrition action plan to meet your specific lifestyle and nutrition goals.

The MyPlate website (<http://www.choosemyplate.gov>) contains nutrition information, healthy eating tips, and ideas for increasing physical activity.



References

Academy of Nutrition and Dietetics. (n.d.). Nutrition Care Manual products. <http://www.nutritioncaremanual.org>

Clark, A., Imran, J., Madni, T., & Wolf, S.E. (2017). Nutrition and metabolism in burn patients. *Burns & Trauma*, 5:11.

Mueller, C., Lord, L., Marian, M., McClave, S., & Miller, S. (Eds.). (2017). *The ASPEN adult nutrition support core curriculum* (3rd ed.). American Society for Parenteral and Enteral Nutrition.

U.S. Department of Agriculture. (n.d.). MyPlate. <http://www.choosemyplate.gov>



Authorship

Healthy Eating After Burn Injury for Adults was developed by Megan Nordlund MS, RD, CD, clinical dietitian, and Nicole S Gibran MD, FACS, UW Medicine Regional Burn Center, Harborview Medical Center, Seattle, Washington, in collaboration with the Model Systems Knowledge Translation Center (MSKTC). It was reviewed and updated by Megan Fobar, MS, RD, CNSC, Haig Yenikomshian, MD, Elizabeth Flores, BA, and Elizabeth Mojarro-Huang, BA, in collaboration with the MSKTC (2023).

Source: The content in this factsheet is based on research and/or professional consensus. This content has been reviewed and approved by experts from the Burn Model System (BMS) centers, funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR). The content of the factsheet has also been reviewed by individuals with burn injury and/or their family members.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this factsheet were originally developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR; grant number 90DP0012) and were updated under an NIDILRR grant (90DPKT0009). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Recommended citation: Fobar, M, Yenikomshian, H., Flores, E., & Mojarro-Huang, E. (2023). *Healthy eating after burn injury—for adults*. Model Systems Knowledge Translation Center (MSKTC).

<https://msktc.org/burn/factsheets/healthy-eating-after-burn-injury-adults>.

Copyright © 2023 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Healthy Eating After Burn Injury—for Kids

November 2023

<https://msktc.org/burn/factsheets>

BURN Factsheet

This factsheet explains the importance of proper nutrition as your child recovers and heals from a burn injury. It describes the nutritional needs during recovery in the hospital and at home, offers tips for a healthy diet for healing, and includes a few kid-friendly recipes.

This factsheet is intended to inform families of children with burn injury about nutrition during hospitalization and after they return home. Your child needs adequate nutrition to grow and develop. Having a burn injury dramatically increases the need for proper nutrition. The larger the burn size, the more nutrition your child needs to heal. A diet high in calories and protein:

- Supports the immune system to decrease risk of infection
- Helps wounds heal faster
- Maintains muscle mass
- Minimizes weight loss to support rehabilitation



How Are Nutrition Needs Determined?

For a child with a burn injury, a dietitian and the medical team decide how much nutrition (calories and protein) your child needs based on their weight, height, age, and burn size. Vitamins and minerals, such as vitamin C, vitamin A, and zinc, are also important for healing and preventing infection. Eating a healthy, well-balanced diet will help your child receive enough nutrients to support wound healing. Your medical team will let you know if your child needs to take any extra supplements.

How Are Nutrition Needs Met While Hospitalized?

Healing from a burn injury requires more calories and protein than any other type of injury. In the hospital, a dietitian makes sure that your child is getting enough nutrients to heal. The dietitian monitors your child's weight, nutrient intakes and outputs, and wound healing.



After a burn injury, it is important to eat a balanced diet that includes a lot of protein, in addition to other foods such as fruits, vegetables, and grains. Your child should eat high-protein foods at every meal and as snacks.

Examples of high-protein foods include:

- Lean meat
- Fish
- Eggs
- Dairy products (milk, yogurt, cheese)
- Beans
- Nuts

The Model Systems Knowledge Translation Center works with Burn Model System (BMS) centers to provide free research-based rehabilitation resources for people living with burn injury (See <https://msktc.org/burn> for more information). This factsheet has been approved by experts from the BMS centers.



Model Systems
Knowledge Translation
Center

Drinking milkshakes or smoothies may help your child to meet their calorie and protein needs. If needed, the medical team may also recommend certain vitamin or mineral supplements.

Your child may need more nutrients than what they get from eating only by mouth. If this is the case, tube feeding can deliver more nutrients. A soft, flexible tube inserted through the nose reaches the stomach to deliver a liquid formula that contains all the nutrients needed for healing. Tube feeding will continue as long as necessary.

What Should Your Child Eat at Home?

After leaving the hospital, you can monitor your child's nutrition status by watching their weight, growth, and wound healing. Take your child to regular doctor appointments where they will chart your child's growth. Remember, your child requires fewer calories than when they were hospitalized. If your child's burn wounds are still open, their diet should include extra protein. As your child continues to heal, their nutrition needs will be like they were before the injury. At the hospital, they likely ate large meals, drank nutrition supplements, and ate a lot of snacks, so your child's appetite may be big when you get home. Now it is important to focus on a balanced diet. Avoid foods with little nutritional value, such as sugary beverages, desserts, candy, fatty meats, and white breads or crackers. Eat more lean meats, whole grains, vegetables, fruits, and dairy. A child with a burn injury needs the nutrients from these foods to continue healing and maintain a healthy weight.



Your child may also need extra vitamins. For example, the health care team may ask your child to take a vitamin D supplement because of extended hospitalization, immobility, and decreased exposure to the sun.

Ask your child's health care provider about exercise to help maintain a healthy weight. Playing and exercising is great for the mind and body. In general, your child can do many of the things that they did before the injury. Be sure to listen to your child's doctor about any limitations. Protect your child's skin from sunlight when they are outside. For more information, refer to the Exercise After Burn Injury Factsheet (https://msktc.org/lib/docs/Factsheets/Burn_Exercise_Fact_Sheet_508.pdf).



Tips for a Well-Balanced Diet at Home

- Offer your child small, frequent meals and snacks. Let him or her decide how much to eat at one time.
- Prepare balanced meals that include all five food groups: fruits, vegetables, grains, protein, and dairy.
- Give your child foods that are high in protein with every meal and snack.
- Have healthy snacks prepared and available in a place where your child can easily find them. For example, consider keeping cheese cubes and apple slices in a container on a low shelf in the refrigerator.
- Make smoothies or milkshakes for your child if they aren't interested in solid food.
- Offer your child water between meals. Sugary beverages such as soda and sports drinks add too many calories with little nutritional value.
- Get creative! Try combining new foods.
- Be a role model: Drink plenty of water, talk about your healthy food choices, and eat with your child.



Ideas for Increasing Protein

When your child eats:	Add or use:
Fruit and vegetable sticks	Peanut butter, almond butter, hummus, or cheese
Whole wheat bread or toast	Peanut butter or melted cheese
Oatmeal	Milk instead of water and add nuts
Crackers or chips	Choose whole grain and eat with peanut butter, cheese, or hummus
Milk	1 cup of regular dry milk powder to 1 quart of milk
Broth-based soups	Cream-based soups
Soups and casseroles	Diced or ground beef, chicken, or turkey
Ice cream or yogurt	Nuts and granola with seeds

It is best to add protein to the diet with whole foods; protein powder supplements are generally not needed.



Kid-Friendly Smoothie Recipes

Monkey Shake (485 calories, 14 g protein)

- 1 banana
- 2 tbsp. peanut butter
- 2 tbsp. chocolate syrup
- ½ cup whole milk

Creamsicle Smoothie (455 calories, 13 g protein)

- 1 cup orange sherbet
- ½ cup whole milk
- 6 oz. vanilla yogurt

Dinosaur Juice (310 calories, 9 g protein)

- ½ cup vanilla ice cream
- 1 or 2 handfuls fresh spinach leaves
- 2 cups frozen fruit: pineapple, mango, or berries
- 2 tbsp. wheat germ or flax seed

Summer Vacation Smoothie (370 calories, 9 g protein)

- ½ cup vanilla ice cream
- 1 package vanilla nutritional powder drink mix
- ½ cup orange juice
- ½ cup frozen fruit (e.g., pineapple, mango, etc.)



Additional Resources

Contact your local burn center and ask for an appointment with a dietitian who will create a nutrition action plan to meet your child's specific lifestyle and nutrition goals.

The MyPlate website (<http://www.choosemyplate.gov/kids>) contains nutrition information, healthy eating tips, and ideas for increasing physical activity.

References

Academy of Nutrition and Dietetics. (n.d.). *Pediatric Nutrition Care Manual products*.
<http://www.nutritioncaremanual.org>

Corkins, M. R. (Ed.). (2015). *The A.S.P.E.N. Pediatric nutrition support core curriculum* (2nd ed.). American Society for Parenteral and Enteral Nutrition.

U.S. Department of Agriculture. (n.d.). *Kids. MyPlate*. <http://www.choosemyplate.gov/kids>



Authorship

Healthy Eating After Burn Injury—For Kids was originally developed in 2016 by Megan Nordlund MS, RD, CD, Nicole S Gibran, MD, FACS, and Maggie Dylewski, PhD, RD, LD, in collaboration with the Model Systems Knowledge Translation Center (MSKTC). It was reviewed and updated by Megan Fobar, MS, RD, CNSC, Haig Yenikomshian, MD, Elizabeth Flores, BA, and Elizabeth Mojarro-Huang, BA, in collaboration with the MSKTC.

Source: The content in this factsheet is based on research and/or professional consensus. This content has been reviewed and approved by experts from the Burn Model System (BMS) centers, funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR). The content of the factsheet has also been reviewed by individuals with burn injury and/or their family members.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this factsheet were originally developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR; grant number 90DP0012) and were updated under an NIDILRR grant (90DPKT0009). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Recommended citation: Fobar, M., Yenikomshian, H., Flores, E., & Mojarro-Huang, E. (2023). *Healthy eating after burn injury—For kids*. Model Systems Knowledge Translation Center (MSKTC).
<https://msktc.org/burn/factsheets/healthy-eating-after-burn-injury-kids>.

Copyright © 2023 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Help Your Child Recover—Build Your Child's Resilience After a Burn Injury

July 2022

<https://msktc.org/burn/factsheets>

BURN Factsheet

This factsheet provides information and tips to help you and your child after a burn injury.

Having a burn injury is a hard experience for children, their parents and guardians, and their families. But it is also an opportunity to build resilience. Resilience is the ability to adapt and overcome challenges. Resilience also refers to the ability of other people and systems to help the child overcome challenges. Children who are resilient often have caring and engaged parents or guardians, a supportive extended family, support from a burn center and their community, and supportive schools. This fact sheet is meant to help parents and guardians understand possible emotional reactions after a burn injury and to provide strategies that promote resilience.




What to Expect After a Burn Injury

Burn injury can be stressful on children and their families. Some children may have dramatic changes in behavior and emotions after a burn. Anxiety, sadness, irritability, and adjustment problems are some of the many changes that children experience and parents often see. For some children, these changes last only a short time, but other children may struggle for weeks or months to adjust to life after a burn. No two children respond the same way. A child's response to a traumatic event can vary based on their age, personality, developmental stage, and the social support they receive. Several reactions are common following a burn injury:

- **Sadness and withdrawal.** Your child may be quiet and may not want to talk to others. They may appear sad, have little interest in activities, or seem more emotional than normal.
- **Regression.** Your child may act younger than his or her age. For example, a preschool-aged child who was toilet-trained may go back to using diapers. An older child may be less independent or suddenly rely heavily on a parent or guardian.
- **Loss of independence.** Your child may be needy and less sure of themselves.
- **Fear.** Your child may be afraid of being alone and may have strong reactions to anything that reminds them of the burn.
- **Separation anxiety.** Your child may get upset about being away from parents or loved ones.
- **Sleeping problems.** Your child may not be able to fall sleep or stay asleep.
- **Nightmares.** Your child may have nightmares.



The Model Systems Knowledge Translation Center works with Burn Model System centers to provide free research-based rehabilitation resources for people living with burn injury (See <https://msktc.org/burn> for more information).



Model Systems
Knowledge Translation
Center

- **Irritability or aggression.** Your child may be angry or moody or lash out.
- **Appetite changes.** Your child may not want to eat or may want to eat too much.
- **Adjustment difficulties.** Your child may have a tough time adjusting to the hospital or to changes that have occurred because of the burn.

These changes in behavior may recur, even after treatment. Some children may have new emotional changes or changes in behavior as new stressors arise.

What Is Acute Stress Disorder and Posttraumatic Stress Disorder?

Some children have strong reactions after a traumatic burn injury, including

- Feeling like the burn is happening all over again,
- Having nightmares,
- Feeling edgy or overly anxious,
- Being overly on guard about getting burned again, and
- Trying to avoid anything that reminds them of the burn.



All these reactions, called acute stress symptoms, are normal in the first three days after the burn injury. The reactions typically go away with time and support. If they don't get better or if they interfere with the child's ability to function, the child may develop acute stress disorder (if occurring within one month of the injury) or posttraumatic stress disorder (PTSD) (if the symptoms persist for months after the injury). The good news is that there are several effective treatments for posttraumatic stress, and most children recover fully. **Parents and siblings may also experience symptoms of PTSD.** To learn more about the symptoms of acute stress disorder and PTSD, please see the following fact sheet: <https://msktc.org/burn/factsheets/PTSD-After-Burn-Injury>.

Building Resilience In Children

Parents and guardians are an important part of a child's emotional recovery. Positive family support is one of the biggest predictors of a child's recovery and adjustment. You can help your child by being present and available, listening to them, and validating his or her concerns. It is also important to model a positive, hopeful, and adaptive way of reacting to challenges. Children look to their parents and guardians for cues on how to react to difficult situations; you can serve as a model for your child as they navigate this difficult time. One of the best ways to model positive coping is to return to a normal routine as soon as possible, and wound care, stretching, etc. can be worked into this routine. Children thrive in environments that are predictable. Having a predictable schedule and routine that closely mirrors what they were doing before the burn is important. This includes returning to school as soon as they can and the parent(s) returning to work as soon as possible after the burn.

Some children may want to talk about the event and/or burn; for these children, it can be helpful to let them talk about their experiences and answer questions. It's also helpful to gently make sure that the child ends their burn story with the understanding that the burn event is over and that they are safe. Other children may not be ready to talk about their burn, and that is okay. There is no need to push your child to talk. Respect how your child responds and watch for signs of anxiety or discomfort. Some children may not be able to communicate their needs verbally, but you can still watch their



behavior for clues of what they need. Reach out to your burn care team or pediatrician who can put you in touch with a mental health professional with training in trauma care if your child:

- Has a change in behavior that lasts for one month after coming home from the hospital. There is a typical period of adjustment after a burn injury. However, if your child's behavior hasn't returned to what it was before the burn within a month after their injury, think about getting help.
- Has behaviors that get in the way of returning to normal routines. For example, your child may be physically ready to return to school, but they may be scared and not want to leave home.

There are several ways to help your child build resilience after a burn:

- Spend time with your child through play, talking, fun activities, and shared activities. Make sure that your child feels loved, safe, and valued.
- Build self-esteem and confidence in your child by allowing them to have as much control and independence as they can handle in a responsible way. Encourage your child to make choices after the burn, even if they are small choices. Encourage your child to problem solve when faced with difficulties. Praise your child when you see them trying to respond well to challenges.
- Model how to respond to challenges in a positive way. It may be helpful to think out loud to help your child understand your thought process. Accept that your child may look differently now. Your child may have a difficult time accepting that they look different from their peers. What your child needs first is acceptance from you—the parents.
- Have the same rules and expectations for your child after the burn as you did for them before the burn. After a burn, parents might have the tendency to go easy on their child or not enforce rules about behavior. This can cause a child to worry. For example, they may start thinking, "I must be really hurt if my mom/dad is letting me get away with this." Your child may not be able to do the same chores that they did before the burn. But you can find appropriate chores that your child can do to contribute to the family.
- Help older children find meaning in what they have been through. Many older children struggle with how much their lives have changed after a burn. But over time, these children may also experience some positive changes, called posttraumatic growth. **Dealing with adversity leads to growth.** Help your child to identify personal strengths and new activities and help them feel hope for the future.
- Connect with family, community, or religious supports. Make sure that your child has support from close family members and loved ones. Consider introducing your child to other children who have had a burn injury. Your area may offer summer camps for children with burn injuries. Ask your local burn provider for a list of these camps.
- Ask if your hospital has school re-entry services. Work with your child's school to make sure that staff are ready to support your child when they return to school.



Parental and Sibling Reactions to a Child's Burn

Parents or guardians and siblings may also have stress symptoms after a child's burn. This is known as secondary traumatic stress. These feelings are expected and common. It is important for parents and guardians to recognize their own emotional reactions to their child's burn injury and to take care of themselves and seek help if the reactions don't go away with time.



Things you can do:

Take time for yourself. Get back into the activities that relax and energize you. Socialize with friends, meditate, take walks, and connect with your spouse. It is important for your child to see that you are getting back into your normal routine.

Make sure that you eat well, get enough sleep, and exercise. It is important for you to be physically and emotionally healthy so that you can be the best support for your child.

Recognize when you feel sad or guilty about your child's injury and recognize when these feelings affect your parenting. For example, if you feel guilty, you might be more lenient and less strict about your child's behavior. You might let them get away with talking to you disrespectfully. You might allow your child to make demands or give them a lot more gifts than you normally would. These changes in your parenting style can have a negative impact on your child. Instead, say, "I love you enough to know that you will get through this and we will be cheering you on." If you always rescue your child from discomfort and frustration, they may not learn how to handle such emotions and challenges. Believe in your child's strength, and they will believe in themselves.



Consider peer support by asking your burn center to introduce you to other parents who have a child with a burn injury. The Phoenix Society trains peer supporters through its SOAR program and offers online support groups for parents of children with burn injuries.

Consider seeking counseling for more support.

Siblings are also affected by the burn injury. Consider the following when supporting siblings:

- Always include siblings in discussions about the burn injury. Never blame siblings for the circumstances of the burn.
- Give extra attention or spend one-on-one time with your other children if you are able.
- Seek out camps or support groups for siblings of survivors of trauma.
- Consider getting siblings involved in caring for your child with a burn. Let siblings help IF THEY WANT TO, but don't force them to help. Encourage siblings to get back to their own activities.
- Consider seeking counseling for siblings if they are struggling.



Stressful life events, such as a burn, will happen. But you can choose to use those events to build resiliency in your child, so that they don't feel like a victim or more vulnerable.



Resources

The National Child Traumatic Stress Network: <https://www.nctsn.org/audiences/families-and-caregivers>

The Phoenix Society: <http://www.phoenix-society.org>

The Knowledge Translation Center Fact Sheet on PTSD: <https://msktc.org/burn/factsheets/PTSD-After-Burn-Injury>



Authorship

Help Your Child Recover—Build Your Child's Resilience After a Burn Injury was developed by Shelley Wiechman, PhD, ABPP; and Brent Smith, PhD, in collaboration with the Model Systems Knowledge Translation Center.

Source: The health information content in this fact sheet is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems Program.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider about specific medical concerns or treatment. The contents of this fact sheet were originally developed under a grant (number H133A110004) from the U.S. Department of Education, National Institute on Disability and Rehabilitation Research. It was updated under the American Institutes for Research Model Systems Knowledge Translation Center, with funding from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant numbers 90DPBU0004, 90DP0082, and 90DPKT0009). NIDILRR is a Center within the Administration for Community Living (ACL), U.S. Department of Health and Human Services (HHS). The contents of this fact sheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Recommended Citation: Wiechman, S. & Smith, B. (2022). *Help your child recover—Build your child's resilience after a burn injury*. Model Systems Knowledge Translation Center (MSKTC). <https://msktc.org/burn/factsheets/Build-Childs-Resilience-After-Burn-Injury>.

Copyright © 2022 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Itchy Skin After Burn Injury

August 2024

<https://msktc.org/burn/factsheets>

BURN Factsheet

This fact sheet explains itchy skin after burn injury, why it occurs, ways to talk about it with your health care team, and strategies for managing it.

Introduction

As skin heals from a burn injury, it may get itchy. Almost everyone recovering from major burns experiences itchy skin—especially on or around the burn, graft, or donor site. The medical term for itchiness is “pruritus” (proo-ri’tus).



Itch intensity (how bad it is) and frequency (how often it occurs) are not necessarily related to the size or depth of your burn injury, but itching may cause you to scratch and open fragile skin that was previously healed. Itching can also make you anxious, which can make itching worse.

Itching can also get in the way of doing everyday activities, including:

- **Sleep.** Itching tends to worsen at night, making it hard to fall asleep and stay asleep.
- **Work or school.** Persistent itching may make it hard to concentrate when performing these activities.
- **Physical activities.** Exercise, sports, and play (for children) may increase itching.
- **Social activities.** The continuous need to scratch an itch may make it uncomfortable to be in social settings.



For many people, itching improves over time. In the meantime, there are treatments to help reduce itching. To find the best treatment for you, talk with your health care team about how bad your itching is and how it is affecting your life.


The Model Systems Knowledge Translation Center works with Burn Model System (BMS) centers to provide free research-based rehabilitation resources for people living with burn injury. (See <https://msktc.org/burn> for more information.) This factsheet has been approved by experts from the BMS centers.

Ways to Describe Your Itching


Your health care team may ask you to describe the **intensity** and **impact** of your itch.

To describe **intensity**, or how strong the itch is, adults are usually asked to rate the itching on a scale of 0 to 10, where 0 is “no itch” and 10 is “worst itch imaginable.” Providers may also ask about how often (hours per day) you itch, the location of the itch, and the impact of the itch on your daily activities.

Children who are unable to describe the details of their itch symptoms in words are often asked to use the Itch Man Scale to describe their itch (see the figure below).¹



Model Systems
Knowledge Translation
Center



¹Source: Itch Man Scale to rate itching intensity in children, designed by Blakeney and Marvin, 2000, at Shriners Hospital for Children. (Reprinted with permission of Shriners Hospital for Children.)

Treatments

Creams and Lotions

- Itching increases with dry skin. If your skin is dry, applying moisturizer may help reduce itching and hydrate the skin. Lotions generally come in a bottle, have a higher water content than creams, and generally soak into the skin faster than creams, but you will need to apply them more often. Creams come in a tube or jar and are generally thicker, so you can apply them less frequently. Apply your moisturizer of choice as often as needed or as instructed by your health care provider.

You should choose the moisturizer that feels best on your skin. Look for unscented or “fragrance-free” moisturizers, as well as moisturizers that do not include alcohol. Products that include alcohol as an ingredient can make your skin dry. (Note: the ingredient “cetyl alcohol” is OK and will not dry out your skin.) For quick and easy tips on selecting a moisturizer, see: [How to Select a Moisturizer for Itchy Skin After Burn Injury](#).



- There are also creams and lotions that have medications in them to help control itching. Some medicated moisturizers may be found over the counter (such as Capsaicin), while others need to be prescribed by your health care provider. Speak with your health care team to see what options may work best for you and your needs.

Bathing

- Bathing may or may not help with itching. Lukewarm water is best and tends to dry out skin the least.
- Bath additives such as oatmeal, powdered milk, or baby oil can be helpful.
- Cooling the skin with water may also provide some temporary relief.



Oral Medications

- Gabapentin, pregabalin, and tricyclic antidepressants** are commonly used oral medications that must be prescribed. They act centrally—in your brain—to help decrease itching.
- Antihistamines** work by blocking histamine, which triggers itching. Common antihistamines that can be purchased over the counter include diphenhydramine (e.g., Benadryl), cetirizine, loratadine, and hydroxyzine.



The correct dosage (how much you should take) varies, so talk with your health care provider before taking any over-the-counter medications.

- **Sleep medications** are used to treat sleeplessness (insomnia). If itching makes it difficult for you to sleep, talk with your health care provider about safe sleep medications that can be helpful to get to sleep and stay asleep.
- **Before taking any medication, talk to your health care provider about your options and potential side effects.** In addition, ask your doctor or health care provider to review all your medications as some medicines can cause itching.

Other Tips

- Pressure garments and other supportive dressings (such as Tubigrip® elasticated tubular bandage, Elastinet®) or tight-fitting sport clothing may help during the first months and up to the first year following a burn injury.
- Skin massage with lotion and/or touching with firm pressure on healed areas can help reduce itching. Strongly patting on the itchy spot rather than scratching is preferable. Make sure you check with your health care provider if your itching remains bothersome.
- Laser therapy is a newer form of treatment that may help with itchy skin. This method provides gradual relief and may require three to six sessions of therapy to get the most benefit. Learn more from our Laser Therapy for Burn Scars factsheet: <https://msktc.org/burn/factsheets/laser-therapy-burn-scars>
- Distractions such as television, games, and low-impact or mild exercise or activities can help take your mind off the itching.
- Short fingernails or sleeping with cotton gloves on, especially for children who tend to scratch at night while asleep, can reduce injuries to fragile skin.
- Avoid the use of fragrances on your skin. Choose unscented laundry detergent and body and bath products (e.g., soaps, lotions, sunscreens).
- Protect your skin from the sun by wearing appropriate clothing (e.g., a hat and long sleeves) and using sunscreen.



For more information

- Infographic: How to Select a Moisturizer for Itchy Skin After Burn Injury (https://bit.ly/Moisturizer_Infographic)
- Infocomic: Itchy Skin After Burn Injury (https://bit.ly/Itchy_Skin_Infocomic)
- The Phoenix Society for Burn Survivors (<http://www.phoenix-society.org/>)
- American Burn Association (<http://www.ameriburn.org/>)
- In addition, you can contact your health care providers about local support groups and other resources.



References

- Carrougner, G. J., Martinez, E. M., McMullen, K. S., Fauerbach, J. A., Holavanahalli, R. K., Herndon, D. N., Wiechman, S. A., Engrav, L. H., & Gibran, N. S. (2013). Pruritus in adult burn survivors: Post-burn prevalence and risk factors associated with increased intensity. *Journal of Burn Care & Research*, 34(1), 94–101.
- Elman, S., Hynan, L. S., Gabriel, V., & Mayo, M. J. (2010). The 5-D itch scale: A new measure of pruritus. *British Journal of Dermatology*, 162(3), 587–593.
- Morris, V., Murphy, L. M., Rosenberg, M., Rosenberg, L., Holzer, C. E. 3rd, & Meyer, W. J. 3rd. (2012). Itch assessment scale for the pediatric burn survivor. *Journal of Burn Care & Research*, 33(3), 419–424.
- Schneider, J. C., Nadler, D. L., Herndon, D. N., Kowalske, K., Matthews, K., Wiechman, S. A., Carrougner, G. J., Gibran, N. S., Meyer, W. J., Sheridan, R. L., & Ryan, C. M. (2015). Pruritus in pediatric burn survivors: Defining the clinical course. *Journal of Burn Care & Research*, 36(1), 151–158.



Authorship

Itchy Skin After Burn Injury was originally developed in 2012 by Gretchen J. Carrougner, RN, MN, and Walter J. Meyer III, MD, in collaboration with the Model Systems Knowledge Translation Center (MSKTC). This factsheet was reviewed and updated in 2016 by Gretchen Carrougner, MN, RN, Northwest Regional Burn Model System; Karen Kowalske, MD, University of Texas Southwestern; Jeffrey C. Schneider, MD; Colleen M. Ryan, MD; and Barbara A. Gilcrest, MD, Boston-Harvard Burn Injury Model System in collaboration with the MSKTC. This factsheet was reviewed and updated in 2024 by Lauren Shepler, MPH; Anupama Mehta, MD; Colleen M. Ryan, MD; and Jeffrey C. Schneider, MD, Boston-Harvard Burn Injury Model System in collaboration with the MSKTC.

Source: The content in this factsheet is based on research and/or professional consensus. This content has been reviewed and approved by experts from the Burn Model System (BMS) centers, funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR). The content of the factsheet has also been reviewed by individuals with burn injury and/or their family members.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider about specific medical concerns or treatment. The contents of this factsheet were developed under a grant (number H133A110004) from the U.S. Department of Education (ED), National Institute on Disability and Rehabilitation Research (NIDRR). It was updated under grants (numbers 90DP0012 and 90DPKT0009) from the NIDILRR. NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, HHS, or ED, and you should not assume endorsement by the federal government.

Recommended citation: Shepler, L., Mehta, A., Ryan, C. M., & Schneider, J. C. (2024). *Itchy skin after burn injury*. Model Systems Knowledge Translation Center. <https://msktc.org/burn/factsheets/itchy-skin-after-burn-injury>.

Copyright © 2024 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Laser Therapy for Burn Scars

August 2024

<https://msktc.org/burn/factsheets>

BURN Factsheet

This factsheet describes the use of laser therapy to treat burn scars. It describes the symptoms that lasers help with as well as any pain and problems associated with laser therapy.

What Is a Laser?

A laser is a device that creates a special, single color of light. Unlike a flashlight, the kind of light a laser creates is focused and full of energy. People use lasers for many things. They use lasers to read barcodes at a store, play music on a CD, measure temperature, and even for medical treatments. Each of these uses requires a specific and different kind of laser. In patients with burns, doctors use lasers to make burn scars look and feel more like normal skin.



What Is a Hypertrophic Scar?

- Any injury to the skin creates a scar. The kind of scar that forms depends on many factors. These include where the injury is on the body, how it gets treated, and the patient's genes. A hypertrophic scar is thick, stiff, and raised above the level of the skin around it. These scars can be mild or severe and may even change how the body moves when they form near joints.
- Burns that are deep, need a long time to heal, or are on parts of the body that move a lot are more likely to develop hypertrophic scars. But these scars can develop anywhere on the body where skin gets injured, including an area where there was a skin graft.
- These scars often appear 2–6 weeks after the skin injury. They continue to develop over the next 6–12 months. These scars often stop growing around 1 year after the injury.
- While growing, hypertrophic scars may be pink, purple, or red. As they get older, they often become dark and brown.
- Hypertrophic scars can look like keloids and their treatments are similar. Unlike keloids, hypertrophic scars do not invade into healthy tissue.
- To learn more about hypertrophic scars, please refer to the *Scar Management After Burn Injury* factsheet at <https://msktc.org/burn/factsheets/scar-management-after-burn-injury>.



The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <https://msktc.org/burn/model-system-centers> for more information). This factsheet has been approved by experts from the BMS centers.

What Are the Symptoms of a Hypertrophic Scar?

People often have the following symptoms:

- **Pain.** People describe this pain as feeling like electricity, stinging, or burning.
- **Itching.** Patients often have an intense desire to scratch these scars, which can create new open wounds that create new scars.
- **A tight feeling.** This often happens when hypertrophic scars develop over parts of the body that move.



Model Systems
Knowledge Translation
Center

Non-Laser Management for Hypertrophic Scars

You can use the following methods with or without laser treatment. Talk to your doctor about what options are best for you.

Types of Lasers for Hypertrophic Scars

There are multiple types of lasers used for scars, but these are the two most common:

- A carbon dioxide (CO2) laser creates very small holes in the thick hypertrophic scar. These holes break up the abnormal scar and give the body a chance to grow a new, softer scar. Because a CO2 laser makes holes in a scar, it can be uncomfortable. As a result, you may need anesthesia.
- A pulsed dye laser (PDL) targets and destroys pink and red colors in burn scars. This lets the scars fade and have a better color match to unburned skin. This process involves less scar damage and is often less painful for patients.



Doctors can use these lasers alone or together to treat hypertrophic burn scars. Their approach depends on what a patient needs.

How Does the Laser Process Work?

Talk to your health care provider about the following:

- Laser treatment for burn scars is noninvasive and usually starts 3 months after the burns have healed.
- For the pulsed dye laser, patients often describe the pain of treatment as feeling like snapping a rubber band against the skin. For the CO2 laser, they describe the pain as feeling like a bee sting. Depending on the size and location of the scar and the age of the patient, patients may need local or general anesthesia for these treatments. Most patients do not need opioid pain medicine after treatment.
- After they treat the scar with a laser, some doctors may apply topical or injectable medicines like steroids.
- Doctors usually cover the treated areas with antibacterial ointment and gauze. Please talk to your doctor about posttreatment care. They can give you instructions on how and when to remove these dressings.
- Treated areas will usually heal in 1–3 days.
- Improvements in pain, itchiness, and tightness are often gradual. Patients may need three to six treatments to see results.
- Laser treatments are well tolerated. Treatment complications include local infections, skin that is dark or light compared to unburned skin, and irritation at the site of treatment.

Is Laser Therapy Covered by Insurance?

- Not all insurance providers cover laser treatment, and they often require an authorization process to approve treatment.
- To get approval from your health insurer, the hypertrophic scars must affect how the body functions. Also, a doctor with experience in burn care must state that laser treatment would help.
- Your health care provider will request approval from your insurance company.



Summary

Hypertrophic burn scars can be painful, tight, and itchy. Standard methods to treat these scars include steroid injections and compression garments. Laser therapy is a new form of treatment that may help. Most people need three to six sessions of laser therapy to get the most benefit. Working with doctors with experience in treating burn scars is the most important step to getting proper treatment with lasers.

References

- Buta, M. R., Hultman, C. S., Bojovic, B., Schulz, J. T., & Goverman, J. M. (2023). Laser therapy of burn scars: CPT coding and reimbursement challenges. *Lasers in Surgery and Medicine*, 55(5), 443. <https://doi.org/10.1002/lsm.23674>
- Hultman, C. S., Friedstat, J. S., Edkins, R. E., Cairns, B. A., & Meyer, A. A. (2014). Laser resurfacing and remodeling of hypertrophic burn scars: The results of a large, prospective, before-after cohort study, with long-term follow-up. *Annals of Surgery*, 260(3), 519–529; discussion 529–532. <https://doi.org/10.1097/SLA.0000000000000893>.
- Won, P., Cooper, M., Gillenwater, T. J., & Yenikomshian, H. A. (2023). Treatment of hypertrophic burn scars with laser therapy: A review of adverse events. *Annals of Plastic Surgery*, 91(6), 715–719. <https://doi.org/10.1097/SAP.0000000000003712>



Authorship

Laser Therapy for Burn Scars was developed by Maxwell Johnson, MD, Jeremy Goverman, MD, and Haig Yenikomshian, MD, in collaboration with the Model Systems Knowledge Translation Center (MSKTC).

Source: The content in this factsheet is based on research and/or professional consensus. This content has been reviewed and approved by experts from the Burn Model System (BMS) centers, funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR).

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider about specific medical concerns or treatment. The contents of this factsheet were developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DPKT0009). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Recommended citation: Johnson, M., Goverman, J., & Yenikomshian, H. (2024). *Laser therapy for burn scars*. Model Systems Knowledge Translation Center. <https://msktc.org/burn/factsheets/laser-therapy-burn-scars>

Copyright © 2024 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Managing Pain After Burn Injury

December 2023

<https://msktc.org/burn/factsheets>

BURN Factsheet

This fact sheet explains how to understand and manage your pain as you recover and heal from a burn injury. It describes different types of pain and medical and behavioral techniques to help you cope.

The Model Systems Knowledge Translation Center works with Burn Model System (BMS) centers to provide free research-based rehabilitation resources for people living with burn injury (See <https://msktc.org/burn> for more information). This factsheet has been approved by experts from the BMS centers.

Introduction

Pain and discomfort are an unfortunate part of burn injury and recovery. Many of our patients tell us that ongoing pain continues to be a problem after discharge from the hospital.



Continued pain can interfere with every aspect of your life, including:

- *Sleep*: pain can make it difficult for you to fall or stay asleep.
- *Ability to work or go to school*: pain can limit your ability to function or concentrate on the job.
- *Mood*: pain can cause depression and anxiety, especially when the pain is severe and lasts a long time.
- *Quality of life*: pain can keep you from being able to enjoy time with loved ones or do activities that are meaningful.
- *Healing*: pain can get in the way of healing if it keeps you from being able to sleep, eat, or exercise enough.

If you are having pain, tell your health care provider.

Things to remember:

- Burn pain is complex and requires careful assessment by your health care provider to find the best treatment.
- Pain management often requires a multidisciplinary approach that includes both medication and non-medication treatments and involves a team of health care providers, such as psychologists, physical therapists, and pharmacists, working with your physician.
- Pain severity is not necessarily related to the size or seriousness of the injury. Small burns can be very painful, while some large burns are not as painful.

Step 1: Understanding Your Pain

There are many different types of burn pain, and each person's pain is unique. Understanding the type, intensity, and duration of your pain is important for getting the best treatment.



Your health care provider will ask you about several types of pain:

- **Acute pain**: short-term intense pain that typically happens during a procedure like dressing changes or physical therapy.



- **Breakthrough pain:** pain that comes and goes throughout the day, often due to wound healing, contractures (tightened skin), or stretching.
- **Resting pain:** also called “background” pain that is almost always present, even at rest.
- **Chronic pain:** ongoing pain that lasts for 6 months or longer after the wound has healed.
- **Neuropathic pain:** pain that is often described as shooting, burning, and can also feel like pins and needles or stabbing. This type of pain is caused by nerves damaged by the burn injury. As these nerves regenerate (regrow), you may experience burning, tingling, or itching.



You might also be asked to describe the pain in the following ways:

- **Intensity:** how strong the pain is, often rated on a scale of 0–10, with 0 as “no pain” and 10 as “worst pain imaginable.”
- **Duration:** how long it lasts (for example— minutes, hours, days, etc.).
- **Timing:** when it gets worse (during the day, night, or during certain activities).
- **Quality:** how the pain feels (for example—stinging, throbbing, itching, aching, shooting).
- **Interference:** how the pain affects your emotions and your ability to do things like work or go to school.

Other important information that can help your health care providers plan the best treatments for your pain include:

- Your experiences with either acute pain or chronic pain before your burn injury.
- Your experiences with poor sleep, depression, or anxiety before or after your burn injury.
- Pain medications, including self-medication and home remedies, you have taken and used in the past.
- How much your pain limits your ability to do certain things.
- Any activities that make your pain worse or better.

Step 2: Treating Your Pain

Medications

- **Over-the-counter** pain medications, such as acetaminophen and nonsteroidal anti-inflammatory drugs (NSAIDs; ibuprofen and naproxen are examples) can be used for long term pain relief. They are not addictive. These medications are effective for treating muscle pain. Use of NSAIDs for long-term pain management may cause serious side effects and should be used only under the supervision of your health care provider.
- **Opioids** are commonly given in the hospital, after discharge from the hospital, and to help with pain from open wounds. Some examples of these medications are morphine, hydrocodone, hydromorphone, oxycodone, and tramadol. Opioids are not effective for chronic burn pain. Side effects, such as constipation and low mood, can also become a problem, and opioids are addictive. For these reasons, your physician will help you taper off opioids when appropriate.
- **Anticonvulsant medications**, such as gabapentin and pregabalin, are useful for managing nerve pain or itching in some situations. These medications work by changing the way the body experiences pain. Anticonvulsant medications can also help with itching.
- **Sleep medications**, such as melatonin, might be used if pain is interfering with sleep. Talk to your physician about sleep hygiene and safe medications for sleep.



- **Antidepressants** can provide pain relief for some people with chronic pain, even if they are not depressed. Antidepressants can also help with sleep. You might talk to your health care provider about trying antidepressants as one way to manage your chronic pain.

Behavioral Approaches

Rarely do medications take away all of the pain. You may also need to use behavioral approaches to help make pain more manageable. A psychologist with expertise in pain management can work with you to find non-medication approaches that can help. These may include:

- **Physical activity** can help manage pain. Although it may seem counterintuitive to increase your physical activity when you are in pain, it is important to remember that not all pain is a signal of harm and needs rest. Research has shown that the more physically active we are, the less pain that we have and the more we are able to do. It is important to establish a regular exercise routine as soon as your doctor says it is safe. This will increase function, decrease pain, and improve your mood and self-esteem. 
- **Relaxation:** a burn injury puts immense stress on the body that continues for many months during the recovery phase. This stress causes muscle tension that can increase pain. Relaxation techniques can be used to lessen the stress placed on your body. Some of these techniques include deep breathing, yoga, and progressive muscle relaxation.
- **Pacing of activities** can help you control your pain. Daily activity and regular exercise are crucial in order to rebuild your strength and stamina and increase your range of motion. But pushing yourself too far can increase your pain.
 - Pace yourself by gradually increasing your physical activity over time. If you are too sore to move comfortably the day after an activity, you have probably pushed yourself too hard.
 - It is best to reduce your activity level until you are more comfortable. This is a difficult balance as burn recovery can be painful, and some pain may be necessary to progress to your previous level of function. Work closely with your physical and occupational therapists to set up an activity program that is best for you.
- **Cognitive (thinking) techniques** use the power of your thoughts to relieve stress. These techniques include a process called “cognitive restructuring,” which helps you change the way you think about your pain and reassure yourself that the pain is temporary and manageable.
- **Mindfulness meditation** has been shown to be a very effective treatment for pain. The technique is easy to learn. There are programs online that can guide you in a mindfulness exercise, or you can work with a mental health provider.
- **Hypnosis** has been shown to be a powerful tool in relieving both acute and chronic pain. A psychologist can teach you how to do self-hypnosis so you can include it in your daily routine. 

Step 3: Coping With Pain

People have different ways of coping with difficult situations or physical discomfort. Your coping “style” can have a large impact on how much pain you feel or how much the pain bothers you.



In any difficult situation, a person reacts by either trying to change the situation, change themselves, or by “giving up.” The first two options are “active” coping styles and are highly effective in managing stress. The third option is much less helpful and often results in withdrawal or symptoms of depression.

Research has shown that it is best to first think about how much of the situation is under your control, and then pick the best active coping style. If the situation is out of your control, changing how you think about and respond to it can be the best coping style. A psychologist can work with you on developing this kind of coping skill.

It is also important to decide which parts of the situation are under your control. For example, you cannot change the fact that you have sustained a burn injury that has resulted in ongoing pain. “Wishing” the injury had not occurred and dwelling on the “what-ifs” won’t help your pain and may lead to feeling more helpless and depressed. However, focusing on the part of the situation that you can control—such as your own rehabilitation, time spent in physical therapy, doing your daily range-of-motion exercises, and following the pain management strategies suggested by your doctor—can be a highly effective coping strategy.

References

- Ratcliff, S. L., Brown, A., Rosenberg, L., Rosenberg, M., Robert, R. S., Cuervo L. J., Villarreal, C., Thomas, C. R., & Meyer 3rd, W. J. (2006). The effectiveness of a pain and anxiety protocol to treat the acute pediatric burn patient. *Burns*, 32(5), 554–562.
- Romanowski, K. S., Carson, J., Pape, K., Bernal, E., Sharar, S., Wiechman, S., Carter, D., Liu, Y. M., Nitzschke, S., Bhalla, P., Litt, J., Przkora, R., Friedman, B., Popiak, S., Jeng, J., Ryan, C. M., & Joe, V. (2020). American Burn Association Guidelines on the Management of Acute Pain in the Adult Burn Patient: A review of the literature, a compilation of expert opinion, and next steps. *Journal of Burn Care and Research*, 41(6), 1129–1151. doi:10.1093/jbcr/iraa119
- Schneider, J. C., Harris, N. L., El Shami, A., Sheridan, R. L., Schulz 3rd, J. T., Bilodeau, M.-L., Ryan, C. M., (2006). A descriptive review of neuropathic-like pain after burn injury. *Journal of Burn Care & Research*, 27(4), 524–528.
- Wiechman Askay, S., Patterson, D. R., Sharar, S. R., Mason, S., & Faber, B. (2009). Pain management in patients with burn injuries. *International Review of Psychiatry*, 21(6), 522–530. doi:10.3109/09540260903343844



Authorship

Managing Pain After Burn Injury was originally developed by Shelley A. Wiechman, PhD and Shawn T. Mason, PhD, in collaboration with the Model Systems Knowledge Translation Center (MSKTC). It was reviewed and updated in 2017 by Shelley A. Wiechman, PhD, Walter J. Meyer, MD, Jeffrey C. Schneider, MD, Karen Kowalske, MD, Kathryn Epperson, BSN, RN, in collaboration with the MSKTC and in 2023 by Shelley A. Wiechman, PhD, Andie Hall, PharmD, BCCCP, Haig Yenikomshian, MD, and Caitlin M Orton, MPH, in collaboration with the MSKTC.

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this factsheet were originally developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR; grant number 90DP0012) and were updated under an NIDILRR grant (90DPKT0009). NIDILRR is a Center within the Administration for Community Living (ACL), U.S. Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Recommended citation: Wiechman, S. A., Hall, A., Yenikomshian, H., & Orton, C. M. (2023). *Managing pain after burn injury*. Model Systems Knowledge Translation Center (MSKTC).
<https://msktc.org/burn/factsheets/managing-pain-after-burn-injury>.

Copyright © 2023 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Outpatient Opioid Management for Adult Burn Survivors

Update for Community Providers

March 2021

| www.msktc.org/burn/factsheets

| BURN Factsheet for Clinicians

INTRODUCTION

People hospitalized for a burn injury often describe it as one of the most painful injuries one can experience. Opioids are frequently prescribed to help alleviate this pain.

Although there is not one standard approach to acute burn pain management, a goal is for the hospitalized patient to be awake, alert, and comfortable.¹ Although unrealistic to expect 'no pain' after a burn injury, significant and enduring in-hospital pain has been shown to be a predictor of poor post-discharge adjustment, well-being and lower health-related quality of life.^{2,3}

However, the opioid crisis in the United States is not going away. Overuse of prescribed opioid medications is increasing and misuse is on the rise.⁴ The Centers for Disease Control and Prevention (CDC) report that in 2017, 47,600 people died from opioid overuse in the United States and opioids were involved in 67.8% of all deaths from drug overdose.⁵

Postdischarge pain management

Between 86%⁶ and 90%⁷ of hospitalized adult burn patients are given opioid prescriptions at discharge. Two weeks after discharge, 90% of patients no longer use opioids,⁶ and most patients no longer use opioids 30 days after discharge.⁷

For those that are discharged from the hospital with an opioid prescription, consider the following opioid prescribing practices:

- Lower daily amounts of prescribed opioids;⁶
- Prescribe only short-acting opioids;⁶
- Educate patients about opioid use;⁶
- Ask patients to complete a risk assessment;⁶
- Check the Prescription Monitoring Program to identify other medications or drugs of concern.⁸

Aim: To provide information about opioid management for adults after they have been discharged from a hospital for treatment of their acute burn injury. This information will help community healthcare providers in their provision of comprehensive care in the setting of a national opioid crisis.

The Burn Model System is sponsored by the National Institute of Disability, Independent Living, and Rehabilitation Research, U.S. Department of Health and Human Services' Administration for Community Living. (See <http://www.msktc.org/burn/model-system-centers> for more information).



Other Options and Factors to Consider:

Scheduled use of nonsteroidal anti-inflammatory agents (NSAIDs) and acetaminophen is an option for non-opioid pain medication. NSAIDs help with pain relief and inhibit cyclooxygenase and prostaglandin synthesis.^a When added to opioids, NSAIDs produce superior pain relief and an opioid-sparing effect (reduced opioid doses without loss of analgesic efficacy) that is associated with a decrease in some opioid-related adverse side effects.⁹ Together, scheduled use of NSAIDs and acetaminophen in outpatient settings may help to reduce opioid use.

Gabapentin may be prescribed to treat neuropathic pain and itching in burn patients. Gabapentanoids (gabapentin and pregabalin) were originally designed as anticonvulsants but have been used to treat chronic neuropathic pain.^b Gabapentin is increasingly used in burn care. However, the data do not support an opioid-sparing effect with the use of gabapentin.^{10,11} Gabapentin remains in the body for 1 or 2 days after stopping use, and withdrawal symptoms can occur a few days later. Tapering is generally not needed for patients who take lower doses—that is, 300 mg/day or less. If tapering is necessary, daily doses should be reduced at a maximum rate of 300 mg total every few days and based on feedback from the patient.



When added to opioids, NSAIDs produce superior pain relief and an opioid-sparing effect that is associated with a decrease in some opioid-related adverse side effects.⁹

Non-pharmacologic or adjunctive treatment strategies may help to relieve pain and itching. Although the majority of the studies cited below were performed with inpatients, findings may be helpful during outpatient recovery and help when opioids are tapered.

- **Scar massage** (with hypertrophic scarring^c): Preliminary evidence suggests that scar massage may be effective in treating pain and itch in persons with hypertrophic scarring¹² and promotes wound desensitization. However, because the evidence is of poor quality, more controlled clinical trials are needed.¹³
- **Cognitive Behavioral Therapy** such as cognitive restructuring, mindfulness, meditation, relaxation, hypnosis, and virtual reality have been used to reduce acute and chronic pain.¹⁴
- **Moisturizer for healed grafts and burn wounds:** Dry, healed burned skin and grafts are associated with increased itch intensity and discomfort. Thus, healed wounds should be kept moisturized with fragrance-free ointments.¹⁵
- **Interactive gaming:** Interactive gaming has been shown to reduce pain during rehabilitation sessions for minor burn injuries.¹⁶
- **Other options to consider:** Acupuncture, acupressure, laser therapy.
- **Assessment for other concerns:** Assessment and treatment of coexisting problems that impact pain is recommended. These may include pruritus, insomnia, post-traumatic stress, depression and anxiety.

^a NSAIDs have an analgesic ceiling and may be associated with platelet dysfunction, gastrointestinal irritation or bleeding, and renal dysfunction. Like NSAIDs, acetaminophen should be administered on a scheduled basis. When added to opioids, acetaminophen produces better analgesia and an opioid-sparing effect associated with a decrease in some opioid-related events, such as nausea/vomiting and sedation. The maximum dosage of acetaminophen for a normal-sized adult is commonly quoted at 4 g/d, but the manufacturer of Tylenol® in the United States has dropped the maximum daily dose to 3 g/d (<https://tylenol.com/safety-dosing/usage/dosage-for-adults>). Acetaminophen is hepatically cleared and should not be used by patients with liver insufficiency.

^b Gabapentanoids are associated with several adverse effects, including sedation, dizziness, and peripheral edema. In elderly patients, these agents should be used with caution or the dose should be decreased. Gabapentanoids are renally excreted; thus, dosage should be lowered in patients with renal dysfunction.

^c A hypertrophic scar is a thickened, raised, red scar that develops where the skin has been injured, as with a burn.

Other Options and Factors to Consider:

Scheduled use of nonsteroidal anti-inflammatory agents (NSAIDs) and acetaminophen is an option for non-opioid pain medication. NSAIDs help with pain relief and inhibit cyclooxygenase and prostaglandin synthesis.^a When added to opioids, NSAIDs produce superior pain relief and an opioid-sparing effect (reduced opioid doses without loss of analgesic efficacy) that is associated with a decrease in some opioid-related adverse side effects.⁹ Together, scheduled use of NSAIDs and acetaminophen in outpatient settings may help to reduce opioid use.

Gabapentin may be prescribed to treat neuropathic pain and itching in burn patients. Gabapentanoids (gabapentin and pregabalin) were originally designed as anticonvulsants but have been used to treat chronic neuropathic pain.^b Gabapentin is increasingly used in burn care. However, the data do not support an opioid-sparing effect with the use of gabapentin.^{10,11} Gabapentin remains in the body for 1 or 2 days after stopping use, and withdrawal symptoms can occur a few days later. Tapering is generally not needed for patients who take lower doses—that is, 300 mg/day or less. If tapering is necessary, daily doses should be reduced at a maximum rate of 300 mg total every few days and based on feedback from the patient.



When added to opioids, NSAIDs produce superior pain relief and an opioid-sparing effect that is associated with a decrease in some opioid-related adverse side effects.⁹

Non-pharmacologic or adjunctive treatment strategies may help to relieve pain and itching. Although the majority of the studies cited below were performed with inpatients, findings may be helpful during outpatient recovery and help when opioids are tapered.

- **Scar massage** (with hypertrophic scarring^c): Preliminary evidence suggests that scar massage may be effective in treating pain and itch in persons with hypertrophic scarring¹² and promotes wound desensitization. However, because the evidence is of poor quality, more controlled clinical trials are needed.¹³
- **Cognitive Behavioral Therapy** such as cognitive restructuring, mindfulness, meditation, relaxation, hypnosis, and virtual reality have been used to reduce acute and chronic pain.¹⁴
- **Moisturizer for healed grafts and burn wounds:** Dry, healed burned skin and grafts are associated with increased itch intensity and discomfort. Thus, healed wounds should be kept moisturized with fragrance-free ointments.¹⁵
- **Interactive gaming:** Interactive gaming has been shown to reduce pain during rehabilitation sessions for minor burn injuries.¹⁶
- **Other options to consider:** Acupuncture, acupressure, laser therapy.
- **Assessment for other concerns:** Assessment and treatment of coexisting problems that impact pain is recommended. These may include pruritus, insomnia, post-traumatic stress, depression and anxiety.

^a NSAIDs have an analgesic ceiling and may be associated with platelet dysfunction, gastrointestinal irritation or bleeding, and renal dysfunction. Like NSAIDs, acetaminophen should be administered on a scheduled basis. When added to opioids, acetaminophen produces better analgesia and an opioid-sparing effect associated with a decrease in some opioid-related events, such as nausea/vomiting and sedation. The maximum dosage of acetaminophen for a normal-sized adult is commonly quoted at 4 g/d, but the manufacturer of Tylenol® in the United States has dropped the maximum daily dose to 3 g/d (<https://tylenol.com/safety-dosing/usage/dosage-for-adults>). Acetaminophen is hepatically cleared and should not be used by patients with liver insufficiency.

^b Gabapentanoids are associated with several adverse effects, including sedation, dizziness, and peripheral edema. In elderly patients, these agents should be used with caution or the dose should be decreased. Gabapentanoids are renally excreted; thus, dosage should be lowered in patients with renal dysfunction.

^c A hypertrophic scar is a thickened, raised, red scar that develops where the skin has been injured, as with a burn.

Other Resources

- Burn-specific factsheets on pain; pain (<https://msktc.org/burn/factsheets/Managing-Pain-After-Burn-Injury>), itching (<https://msktc.org/burn/factsheets/Itchy-Skin-After-Burn-Injury>), and Sun Exposure (<https://msktc.org/burn/factsheets/sun-protection-after-burn-injury>).

References

1. Faucher L, Furukawa K. Practice guidelines for the management of pain. *J Burn Care Res.* 2006;27(5):659-668.
2. Patterson DR, Tininenko J, Ptacek JT. Pain during burn hospitalization predicts long-term outcome. *J Burn Care Res.* 2006;27(5):719-726.
3. Spronk I, et al. Predictors of health-related quality of life after burn injuries: a systematic review. *Crit Care.* 2018;22.
4. National Institute on Drug Abuse. Overdose death rates. <https://www.drugabuse.gov/related-topics/trends-statistics/overdose-death-rates>. Accessed 08-28-2019, Revised January 2019.
5. Centers for Disease Control and Prevention. <https://www.cdc.gov/drugoverdose/data/statedeaths.html>. Accessed 08-28-2019.
6. Wibbenmeyer L, et al. An evaluation of discharge opioid prescribing practices in a burn population. *J Burn Care Res.* 2015;36(2):329-335.
7. Yenikomshian HA, et al. Outpatient opioid use of burn patients: a retrospective review. *Burns.* 2019; 45(8):1737-1742.
8. WA State Department-of-Health. Prescription Opioids for Acute Pain. 2019. Document # DOH Pub 631-077.
9. Richards D. The Oxford Pain Group League table of analgesic efficacy. *Evid Based Dent.* 2004;5(1):22-23.
10. Kneib CJ, Sibbett SH, Carrougner GJ, et al. The effects of early neuropathic pain control with gabapentin on long-term chronic pain and itch in burn patients. *J Burn Care Res.* 2019;40(4):457-463.
11. Wibbenmeyer L, Eid A, Liao J, et al. Gabapentin is ineffective as an analgesic adjunct in the immediate postburn period. *J Burn Care Res.* 2014;35(2):136-142.
12. Cho YS, et al. The effect of burn rehabilitation massage therapy on hypertrophic scar after burn: a randomized controlled trial. *Burns.* 2014;40(8):1513-1520.
13. Ault P, Plaza A, Paratz J. Scar massage for hypertrophic burns scarring—a systematic review. *Burns.* 2018; 4(1):24-38.
14. Scheffler M, Koranyi S, Meissner W. Efficacy of non-pharmacological interventions for procedural pain relief in adults undergoing burn wound care: a systematic review and meta-analysis of randomized controlled trials. *Burns.* 2018;44:1709-1720.
15. Carrougner GJ, Martinez EM, McMullen KS, et al. Pruritus in adult burn survivors: postburn prevalence and risk factors associated with increased intensity. *J Burn Care Res.* 2013;34(1):94-101.
16. Parker M, Delahunty B, Heberlein N. Interactive gaming consoles reduced pain during acute minor burn rehabilitation: a randomized, pilot trial. *Burns.* 2016;42:91-96.
17. Lawrence R, Mogford D, Colvin L. Systematic review to determine which validated measurement tools can be used to assess risk of problematic analgesic use in patients with chronic pain. *Br J Anaesth.* 2017; 119(6):1092-1109.
18. National Institute on Drug Abuse. <https://www.drugabuse.gov/sites/default/files/files/OpioidRiskTool.pdf>. Accessed 12-16-2019.
19. The Opioid Therapy for Chronic Pain Work Group. VA/DoD Clinical Practice Guideline for Opioid Therapy for Chronic Pain (Version 3.01, 2017). <https://www.healthquality.va.gov/guidelines/Pain/cot/VADoDOTCPG022717.pdf>. Accessed 08-28-2019.

Authorship

Outpatient Pain Management for Burn Survivors—Update for Community Providers was developed by Gretchen J. Carrougner, MN, RN (Northwest Regional Burn Model System); Samuel P. Mandell, MD, MPH, FACS (University of Washington Medicine Regional Burn Center, Seattle, WA); and Katherine D. Travnick, MD (Pain Institute of Nevada, Las Vegas, NV) in collaboration with the Model Systems Knowledge Translation Center.

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Model Systems.

Disclaimer: The contents of this factsheet were developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, HHS, and you should not assume endorsement by the Federal Government.

Copyright © 2021 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.

Post-Traumatic Stress Disorder (PTSD) After Burn Injury

May 2023

<https://msktc.org/burn/factsheets>

BURN Factsheet

This factsheet explains post-traumatic stress disorder (PTSD) after burn injury and common symptoms that may occur.

What Is PTSD?

Mental and physical reactions are common after a traumatic event, like a burn injury. Reactions that occur soon after a traumatic event are called acute stress symptoms. Symptoms of acute or post-traumatic stress are common responses to a burn injury. Think of it as a way for the brain to let the body know it needs to stay safe—these symptoms can actually protect us. Symptoms can develop after experiencing or witnessing a traumatic event (e.g., as a bystander or first responder) or by learning of a bad injury involving a loved one. If symptoms of post-traumatic stress do not go away, they can cause more problems and become Post-Traumatic Stress Disorder (PTSD). PTSD can affect adults and children.



What Are the Symptoms of PTSD?

- Nightmares related to trauma that disrupt sleep or cause restless sleep. Children may not remember the content of the nightmares but remember that the nightmares were frightening.
- Feeling like you're in a daze or detached, or that the burn is not real.
- Intrusive thoughts of the burn injury or feeling as though it's happening again (flashbacks).
- Children may engage in repetitive play in which they "act out" the traumatic event.
- Avoiding reminders of the injury, such as talking about it, watching similar events on TV, or not wanting to return to the place where the trauma occurred.
- Being very easily startled or feeling constantly on guard. New fears may develop, such as fear of the dark, crowds, or elevators. Some people will develop anxiety or a feeling of being on edge or not being safe.
- Negative changes in mood or cognition, such as being irritable or not being able to concentrate.

The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <https://msktc.org/burn/model-system-centers> for more information).

Who Gets PTSD?

About 90% of both adults and children with burn injuries report at least one symptom of acute stress right after the traumatic event but only about 30% of people with these symptoms develop PTSD. PTSD is more likely to occur if the burn injury is an assault or a repeated trauma (such as ongoing abuse).



Burn survivors most at risk for PTSD are those with a history of anxiety disorders (generalized anxiety, panic disorder) or depression. Burn survivors who have a history of traumatic events and past PTSD are also at risk for developing PTSD from the current burn injury. In fact, a current burn injury can trigger nightmares and flashbacks from a past traumatic event. Burn survivors with high levels of pain and anxiety while in the hospital may be more at risk of developing PTSD after discharge. Caregivers and the burn team should effectively manage pain and anxiety from a burn wound to avoid future distress. Children are more at risk of developing PTSD if their parents are experiencing such symptoms.



It is also important to know that some burn survivors may have symptoms of PTSD after the traumatic event, experience a lot of emotional struggles, and then find a new appreciation of life and strength that is even better than before the injury. This is called post-traumatic growth (PTG). People are more likely to have PTG when they use humor, gratefulness, determination, and support from other people.

What Should I Do if I Have Symptoms of PTSD?

For most people, symptoms of post-traumatic stress usually go away over time. After a burn injury, positive social support can be very important. Connecting with friends or family who can provide positive support (no criticism or trivializing your experience) has been shown to be helpful. You may find it useful to get information and details about the trauma. Only talk about the details when you feel ready to do so. Retelling the details of the trauma can actually be re-traumatizing and you can tell friends and strangers who ask that you do not wish to talk about it. However, talking about your emotions is important. Let caregivers and the burn team know if you're worried about how you're coping with the burn injury.

Most early traumatic stress reactions get better over time as you return to a normal routine and start to develop ways to cope. For some people, early reactions may get worse. After the burn injury heals, the emotional trauma may not go away. The intensity and length of PTSD symptoms are not related to the severity of the injury. Seek treatment if symptoms of PTSD last for weeks or months and/or start to interfere with your quality of life.

Very good treatments are available for PTSD and it can go away. The most effective treatments involve a type of counseling called cognitive behavioral therapy (this may also be called Trauma-Focused Cognitive Behavioral Therapy). This is a short-term treatment in which you learn ways to relax and manage anxiety, to gradually be exposed to aspects of the traumatic event, and to think differently about the event and how it affected you. Your doctor may prescribe medicine for a short time to help with anxiety, depression, or sleep problems. While awaiting treatment, you can do several things to treat each symptom:

- **Nightmares.** Wake up and remind yourself of where you are. Re assure yourself that you are safe and go right back to sleep. Try not to get up, turn on the TV, or get something to eat—these will keep you awake longer. Try changing the end of your nightmare by imagining a better outcome.
- **Flashbacks.** Try to stop flashbacks right away by imagining a stop sign or a red light and thinking about something more pleasant. It is important to realize that you have control over what you think about.
- **Anxiety and arousal.** Practice a relaxation technique to reduce your anxiety. Deep or rhythmic breathing, mindfulness meditation, muscle relaxation, yoga, and exercise can all be helpful.



- **Avoidance.** If reminders of the trauma are interfering with your normal routine—such as not wanting to return to work if that is where the accident occurred, or not wanting to go outside near the gas grill—then consider a *fear hierarchy*. A fear hierarchy is a list of steps that slowly expose you to the normal things. For example, consider going back to the workplace for a very short period of time for a casual visit, such as having lunch with a coworker. Then, gradually increase the time you spend in the workplace until you can be near the accident site for an entire shift. Practice relaxation techniques as you work through the fear hierarchy. You may need professional assistance to help with this.

Always talk to the burn team about PTSD symptoms that you may be having. They can help you to find a mental health provider for treatment.

Additional Resources

- National Child Traumatic Stress Network: <http://www.nctsn.org>
- National Institute of Mental Health: <http://www.nimh.nih.gov/health/topics/post-traumatic-stress-disorder-ptsd/index.shtml>

References

- Hobbs, K. (2015). Which factors influence the development of post-traumatic stress disorder in patients with burn injuries? A systematic review of the literature. *Burns*, 41(3), 421–430.
- Hokes, K. E., & Adams, L. M. (2022). The longitudinal development of posttraumatic growth among U.S. adult burn injury survivors. *Rehabilitation Psychology*, 67(3), 369–380.
- Giannoni-Pastor, A., Eiroa-Orosa, F. J., Fidel Kinori, S. G., Arguello, J. M., & Casas, M. (2016). Prevalence and predictors of posttraumatic stress symptomatology among burn survivors: A systematic review and meta-analysis. *Journal of Burn Care & Research*, 37(1), e79–e89.
- Martin, L., Rea, S., & Wood, F. (2021). A quantitative analysis of the relationship between posttraumatic growth, depression and coping styles after burn. *Burns*, 47(8), 1748–1755.
- Van Loey, N., & Van Son, M. J. M. (2003). Psychopathology and psychological problems in patients with burn scars. *American Journal of Clinical Dermatology*, 4(4), 245–272.



Authorship

Post-Traumatic Stress Disorder (PTSD) After Burn Injury was developed by Shelley A. Wiechman, Ph.D., ABPP in collaboration with the Model Systems Knowledge Translation Center (MSKTC). Revision by Kimberly Roaten, Ph.D., Shelley A. Wiechman, Ph.D., ABPP and Caitlin Orton, MPH, in collaboration with the MSKTC.

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this factsheet were developed under a grant (number 90DP0082) from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR) and were updated under a NIDILRR grant (90DPKT0009). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS and you should not assume endorsement by the federal government.

Recommended Citation: Wiechman, S. A., & Orton, C. (2023). *Post-Traumatic Stress Disorder (PTSD) After Burn Injury*. Model Systems Knowledge Translation Center (MSKTC).
<https://msktc.org/burn/factsheets/post-traumatic-stress-disorder-ptsd>.

Copyright © 2023 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Psychological Distress after Burn Injury

November 2016

www.msktc.org/burn/factsheets

BURN Factsheet

This factsheet explains some emotional recovery challenges that burn survivors may face after a major burn injury and treatment options.

Introduction

The purpose of this factsheet is to describe some of the emotional recovery challenges that patients may face after sustaining a major burn injury. It is important to point out that most burn survivors do very well psychologically after surviving a burn injury. However, even though we are optimistic about long term recovery, burn survivors may face several challenges in the aftermath of their injury. Whereas each individual experiences psychological distress differently, people with burn injuries often report:

- Feeling...
 - Sad, anxious or irritable
 - Helpless
 - Hopeless
 - Upset about depending on other people for assistance
 - Distant from family, friends or the general public
 - Alone
 - As if the injury was happening again or reliving it
 - A physical reaction (e.g. heart pounding, trouble breathing, or sweating) when something reminds you of the injury
 - Jumpy or easily startled
 - “Super alert” or watchful and on guard
- Difficulty falling asleep due to thoughts like:
 - “I worry about bad things that might happen”
 - “I keep thinking about the way I was injured”
- Difficulty finding enjoyment in things that used to give pleasure
- Difficulty staying asleep
- Avoiding situations that remind you of the accident
- Avoiding thinking or talking about the injury and how it occurred

While in the hospital, survivors may find they have a lot of time to focus on their burn injury. Many people report having psychological distress several days or a few weeks after they were injured. For most, periods of distress become less frequent and less upsetting after a couple of weeks to a couple of months. However, problems that continue for more than a month or two, or thoughts about wanting to die or hurt oneself, indicates a need to seek treatment.

The Burn Model System is sponsored by the National Institute of Disability, Independent Living, and Rehabilitation Research, U.S. Department of Health and Human Services' Administration for Community Living. (See <http://www.msktc.org/burn/model-system-centers> for more information).



Model Systems
Knowledge Translation
Center

Causes of psychological distress after burn injuries

Major burn injuries can be extremely upsetting and distressing for the survivor, as well as for family members, and friends. Depression or anxiety before the burn injury increases the chances of those symptoms after the injury. Common causes of distress include:

- Thinking about the event itself – both in sustaining the injury and/or witnessing others who were also seriously injured or died
- Worries about the future
- Concerns about finances and the impact the injury has had on family members
- Changes in appearance because of scars and contractures
- Physical discomfort
- Pain while the wound is still healing (especially during the repeated dressing changes) and pain that continues for months afterward as nerves are healing
- Itching
- Difficulty adhering to range of motion and Physical Therapy exercises
- Limitations in physical abilities
- Loss of independence
- Difficulty in returning to work or school
- Loss of property, residence, pets, etc.
- Interruption of daily life activities and roles
- Challenges with sexual interests and intimacy

Effects of psychological distress on health and recovery

Psychological distress has been shown to affect the way the mind works (e.g., poor memory, short attention span) and the ways the body functions (e.g., immune system, digestion). Distress can also worsen other medical conditions (e.g., blood pressure, glucose control) and can interfere with recovery from the burn in many ways, such as:

- Making pain and itching feel even worse
- Reducing your effort and persistence in participating in rehabilitation therapies and wound care
- Making communication with burn team members difficult
- Reducing your interest and pleasure in daily activities
- Disrupting sleep
- Causing tension in interpersonal relationships

Treatment options

It is critical to seek emotional support from professionals and other survivors to help with your psychological distress. There is a caring community that understands your experience. Always let your burn team know about challenges in your emotional recovery.

- Keep connected with friends and family and ask for support
- Take one step at a time during the recovery process. Acceptance of your injury and the changes in your life take time, and recovery (psychological and physical) can proceed at a slow pace
- Get sufficient sleep and eat healthy foods
- Try to avoid napping during the day so that you will sleep better at night
- Avoid tobacco, illicit drugs and/or excessive use of alcohol because they can lead to low mood and increase anxiety
- Stay focused on tasks that you can do rather than those things that are no longer possible because of your injury
- Stay active and exercise regularly
- Return to a normal routine as soon as possible. Get up, get dressed, groom yourself and get out of the house every day to avoid feeling more depressed
- Engage in one positive, pleasant activity every day
- As soon as you are medically cleared, get back to doing the things you did before the injury like going to work or school and doing chores around the house

Peer Support

- Seek support from professionally led support groups. Ask your health care provider how to locate the right group for you
- Seek support from other survivors. You can find survivors who have been trained by the Phoenix Society's SOAR program to provide peer support (www.phoenix-society.org/resources)
- Participate in regularly scheduled peer support on-line discussions offered through the Phoenix Society (www.phoenix-society.org/chat)

Psychotherapy

Mental health professionals are trained in methods for assessing and treating psychological distress.

Professional help is particularly important if the distress is severe and interferes with things that are important to you.

There are many health care practitioners such as psychiatrists, psychologists, social workers, and pastoral counselors that can help. It is best to work with a mental health professional who has experience in treating people with severe injuries and expertise in treating the problems you may be experiencing (e.g., body image, social discomfort, post-traumatic stress disorder (PTSD)).

Here are a couple of the effective methods that health care providers may use to help reduce your distress:

- Cognitive Behavioral Therapy (CBT). Ask your health care provider for more information
- Stress Management: Learning practices like deep breathing, meditation, or staying “present in the moment”
- Coping strategies such as active problem solving
- Communication and social skills: For example, changes in appearance may require that you learn new skills for managing distress that may arise when other people ask about your accident or when they react to changes in your appearance due to your injury

Medications

Consult your primary care doctor or the burn care team to determine if medications may best address your symptoms. Certain medications have been shown to help psychological distress for disorders such as:

- Depression (e.g. low mood, low energy, suicidal ideation, isolation, guilt, irritability towards self and others, loss of interest in things you used to enjoy)
- Anxiety (e.g., worry, recurring and disturbing memories)
- Sleep (e.g., nightmares, difficulty relaxing)

For more information and resources

The Phoenix Society for Burn Survivors: www.phoenix-society.org/

Changing Faces: www.changingfaces.org.uk/

American Burn Association: http://www.ameriburn.org/resources_links.php

Face IT Online: www.faceitonline.org.uk

MSKTC NIDILRR Factsheets: www.msktc.org/burn/factsheets

National Center for PTSD: www.ptsd.va.gov/PTSD/public/treatment/therapy-med/index.asp

Anxiety and Depression Association of America: www.adaa.org/

References

Edwards RR, Smith MT, Klick B, Magyar-Russell G, Haythornthwaite JA, Holavanahalli R, Patterson DR, Blakeney P, Lezotte D, McKibben J, Fauerbach JA. Symptoms of depression and anxiety as unique predictors of pain-related outcomes following burn injury. *Annals of Behavioral Medicine* 2007; 34(3): 312-322.



Fauerbach JA, McKibben J, Bienvenu OJ, Magyar-Russell G, Smith MT, Holavanahalli R, Patterson DR, Wiechman SA, Blakeney P, Lezotte D. Psychological Distress Following Major Burn Injury. *Psychosomatic Medicine* 2007; 69: 473-482.

Logsetty, S., et al. (2016) Mental health outcomes of burn: A longitudinal population-based study of adults hospitalized for burns. *Burns* 42(4): 738-744.

Mason, S.T., Corry, N., Gould, N., Amoyal, N., Gabriel, V., Wiechman Askay, S., Holavanahalli, R., Banks, S., Arceneaux, L. L., Fauerbach, J.A. (2010) Growth Trajectories of Distress in Burn Patients. *Journal of Burn Care Research* 31(1): 64-72.

Mason, S.T., Fauerbach JA, Haythornthwaite, J. Assessment of Acute Pain, Pain Relief and Pain Satisfaction. Chapter 41. in D.C.Turk and R. Melzack (Eds). *Handbook of Pain Assessment: Third Edition* 2010, Guilford Press: New York, NY.

Reeve J, James F, McNeill R. Providing psychosocial and physical rehabilitation advice for patients with burns. *J Adv Nurs*. 2009 May;65(5): 1039-43.

Smith MT, Klick B, Kozachik S, Edwards RR, Holavanahalli R, Wiechman S, Blakeney P, Lezotte D, Fauerbach JA. Sleep onset insomnia symptoms during hospitalization for major burn injury predict chronic pain. *Pain*. 2008 Sep 15; 138(3): 497-506.

Authorship

Psychological Distress was developed by James Fauerbach, Ph.D., Shelley Wiechman, Ph.D., and Shawn Mason, Ph.D. in collaboration with the Model Systems Knowledge Translation Center.


Factsheet Update

This factsheet was reviewed and updated by Shelley Wiechman, PhD, Simrun Kalra, MD, Kimberly Roaten, PhD, Atilla Ceranoglu, MD, and Donna Digioia, MD. The review and update is supported by the American Institutes for Research Model Systems Knowledge Translation Center.

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. This publication was produced by the Burn Model Systems in collaboration with the University of Washington Model Systems Knowledge Translation Center with funding from the National Institute on Disability and Rehabilitation Research in the U.S. Department of Education, grant no. H133A060070. It was updated under the American Institutes for Research Model Systems Knowledge Translation Center, with funding from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this fact sheet do not necessarily represent the policy of the U.S. Department of Education or the U.S. Department of Health and Human Services, and you should not assume endorsement by the federal government.

Copyright © 2016 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Scar Management After Burn Injury

August 2024

<https://msktc.org/burn/factsheets>

BURN Factsheet

This factsheet explains scarring after burn injury, including different types of scars and information about how to manage them.

Scars occur after burn injury. Some scars may impact a burn survivor both physically and mentally after their initial burn injury has healed. It is difficult to predict who will develop scarring, though scarring is related to skin tone and texture prior to burn injury, as well as the depth and location of the burn.



Why Do Scars Form?

Scars form when the dermal or lower layer of the skin has been damaged. The body forms a protein called collagen to help heal the damaged skin. Normally the skin collagen fibers are laid down in a very organized manner, but in scars, these fibers are created in a very disorganized manner, which can give the new skin/scar a different texture and appearance.

How Long Does it Take Scars to Mature?

Scars will “mature” or change over time and can limit a survivor’s ability to function, as well as affect their body image. Scarring usually develops within the first few months after the burn, peaks around 6 months, and may improve or mature in 12–18 months.

Research shows that less severe burns that heal in less than 14 days generally have no scarring. Burns that take longer to heal are at higher risk for problematic scarring and may require further treatment, including reconstructive surgery. The scar maturation process can often seem long and frustrating for a burn survivor and their family. Physically, one might notice their scars:

- Can be dry, leading to cracking or breakdowns of the skin.
- Are more sensitive to touch and their physical environment, for example, exposure to the sun and certain bath or body products.
- Can lead to symptoms such as neuropathic pain or nerve pain, as well as a pins and needle sensation in the scar.

As scars mature, they fade in color and become flatter, softer, and generally less sensitive. People with visible scars may feel self-conscious and avoid social situations. This can lead to isolation, depression, and lower quality of life. If you have concerns or questions about your healing process, treatments, or mental and physical health, contact your health care providers.



The Model Systems Knowledge Translation Center works with Burn Model System (BMS) centers to provide free, research-based rehabilitation resources for people living with burn injury. (See <https://msktc.org/burn> for more information.) This factsheet has been approved by experts from the BMS centers.



Model Systems
Knowledge Translation
Center

Caring for Your Scars

Moisturize

It is important to keep your scars clean and well moisturized for the best outcome. Massage your scars with lotion to keep them moist, make them less sensitive, decrease itching, and make stretching easier. This may also prevent skin breakdown.

- Moisturizers can be applied to all healed areas frequently throughout the day.
- Moisturizers should be applied in thin layers and massaged in gently while the scars are more fragile. As your scars mature, you can begin to add more pressure to help your scars loosen so that they are not so stiff.
- Hot showers remove the natural oils from the skin. Therefore, it is essential to re-moisturize well after showers. Bathing may feel good but also removes the natural oils. Adding baby oil to the water may alleviate some of this effect.

When picking a moisturizer, be sure to avoid products with fragrances or alcohol. You should never use mineral oil, petroleum jelly, or antibiotic ointments to moisturize your skin. These can lead to allergic reactions and skin breakdown. Do not use antibiotic ointment to lubricate after the wound is healed. Check out the below resources for more tips on picking a moisturizer and itchy skin after burn injury. You should also talk to your doctor about what treatments or medications might be right for you.

- **Factsheet:** [Itchy Skin after Burn Injury \(https://bit.ly/Itchy-Skin-Factsheet\)](https://bit.ly/Itchy-Skin-Factsheet)
- **Infographic:** [How to Select a Moisturizer for Itchy Skin After Burn Injury \(https://bit.ly/Moisturizer-Infographic\)](https://bit.ly/Moisturizer-Infographic)

Sun Exposure

You should avoid exposing your scars to sunlight. Scars that are discolored and have not matured burn easily. If you go out in the sun, we recommend

- Planning activities in the early morning or late evening when the sun is the least intense.
- Applying a broad-spectrum sunscreen with an SPF level greater than 30 frequently to maintain protection.
- Re-applying sunscreen every 2 hours.
- For more information, check out our factsheet: [Sun Protection After a Burn Injury \(https://bit.ly/Sun-Protection-Factsheet\)](https://bit.ly/Sun-Protection-Factsheet)

Contractures

Scars across joints can lead to contractures and can affect your ability to move and take care of yourself. If your contractures involve your legs, you may have difficulty squatting, sitting, walking, or climbing stairs. If your contractures involve your trunk and arms, you may have difficulty with grooming, eating, dressing, and bathing as well as working with your hands. Some contractures are unavoidable, but many can be prevented with active involvement in your rehabilitation program. Here are a few reminders:

- Stretching should be performed a minimum of five to six times per day. To make stretching easier, first moisturize your scars with a moisturizer recommended by your doctor.
- Your therapist may make a cast or splint to help position your scar in a stretched position. It is important that you wear the cast or splint as prescribed and tell your therapist if it becomes painful or causes skin irritation.



- Do as much for yourself as possible such as getting dressed and self-grooming. It may take longer than you are used to, but movement and activity will improve your ability to move and take care of yourself.

What Are Hypertrophic Burn Scars?

Hypertrophic burn scars are raised scars in the area of the original burn and are the most common complication of a burn injury. Hypertrophic scars generally develop from more severe burn injuries and:

- Stay within the area of the original burn injury.
- Develop within the first few months after injury.
- Often have a deep red to purple color and are raised above the surface of the skin.
- Can be warm to the touch, very sensitive, and itchy.
- Are more prominent and noticeable around joints where skin tension and movement are high.

What Are Keloid Burn Scars?


Keloid burn scars are very similar to hypertrophic scars, with the major difference being keloid scars do not stay within the area of the original burn injury.



Treatment of Hypertrophic and Keloid Burn Scars

You will need to work closely with your doctor and therapy team to make sure your scars mature as completely as possible. A strong commitment is required from you and your family member to follow through with the treatment plan to ensure the best scar result. No single treatment is ideal for treating hypertrophic scars.

- You and your medical team may decide to use **pressure garments**. They can decrease itching and protect the skin from injury. Some burn survivors also feel that the pressure garments look better than the scars themselves.
- **Gel sheets** are pieces of thin, flexible material that are placed over the scars and may decrease itching and dryness. They are generally durable and comfortable to wear. They can be worn alone or underneath pressure garments, splints, or casts. Some people find that they are sensitive to silicone, so check your skin frequently for irritation or rashes.
- Your therapist may recommend **custom-made inserts** to be worn under gloves, compression bandages, or custom garments to increase pressure on the scar and improve maturation. These inserts can be made from a variety of substances, ranging from soft foam to a rubber consistency.
- **Massage can help soften and desensitize the scar.** When combined with stretching, massage can make the scar looser, softer, and more comfortable. Talk to your therapist to learn about specific massage techniques.
- **Surgical treatment**, including laser treatment, may be an option if scarring prevents you from performing certain activities. It is important to stay in contact with your treating burn physician for evaluation.



For more information regarding compression garments, wound care, and scar management products, please contact your doctor or therapist so that they can make recommendations based on your specific needs.



What Can You Do?

- Be actively involved in your recovery by asking questions and participating in decision making about your care. Take a list of questions or concerns to your medical appointments for your health care provider to address.
- Keep up your exercise program as recommended by your doctor.
- It is important to follow your providers' instructions for using pressure garments, inserts, splints, or gel sheets. If they do not fit properly or if they cause problems such as pressure or skin breakdown, let your health care provider know right away.



Authorship

Scar Management After Burn Injury was originally developed in 2016 by Sandra Hall, PT, Karen Kowalske, MD, and Radha Holavanahalli, PhD, in collaboration with the Model Systems Knowledge Translation Center (MSKTC). This factsheet was reviewed and updated in 2024 by Lauren J. Shepler, MPH, Huan Deng, PhD, Jonathan S. Friedstat, MD, and Jeffrey C. Schneider, MD, Boston-Harvard Burn Injury Model System in collaboration with the MSKTC.

Source: The content in this factsheet is based on research and/or professional consensus. This content has been reviewed and approved by experts from the Burn Model System (BMS) centers, funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR). The content of the factsheet has also been reviewed by individuals with burn injury and/or their family members.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this factsheet were originally developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR; grant number 90DP0012) and were updated under an NIDILRR grant (90DPKT0009). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Recommended citation: Shepler, L. J., Deng, H., Friedstat, J. S., & Schneider, J. C. *Scar management after burn injury*. Model Systems Knowledge Translation Center (MSKTC). <https://msktc.org/burn/factsheets/scar-management-after-burn-injury>

Copyright © 2024 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Sexuality and Intimacy After Burn Injury

March 2021

www.msktc.org/burn/factsheets

BURN Factsheet

This factsheet explains how a burn injury may affect sexuality and intimacy.

Introduction

A burn injury can change the way your body looks, feels, and functions. In some cases, these changes in your appearance may be permanent. After a burn injury, the initial focus is on addressing your immediate, physical needs for survival. Once these needs are met and you are on a path to recovery, issues such as sexuality or intimacy may come up.



What Is Sexuality and Intimacy?

Sex and sexuality are important parts of everyday life. Sex refers to biological and physiological characteristics of your body, and is often used to refer to sexual intercourse. Sexuality refers to how you express yourself, your gender identity (the gender you feel you are), and your sexual identity (your desire for the opposite sex, same sex, or both).

Intimacy is a way to connect closely with oneself or another person on an emotional or physical level. Intimacy with oneself includes, for example, getting comfortable with one's own body via masturbation. You may have different types of intimate relationships; not all intimate relationships are sexual.

Sexuality After Burn Injury

People have different sexual desires and use different ways to express them. Defining what is typical or common is hard. It is whatever gives you and your partner pleasure. It is important to have an open line of communication with your partner to deal with your sexuality.



It is normal to have doubts and fears during this time of healing. These feelings are a natural response to returning to a life of sexuality and intimacy. You are not the only one to feel this way after a burn injury. Talking about body image, sexuality, and intimacy can be hard to do. Sexuality and intimacy are possible regardless of your burn injury.

This factsheet includes information on how changes in your body after a burn injury may affect sexuality and intimacy, suggestions for how to adjust to the changes, and other resources that may help you.

Physical Effects of Burn Injury on Sexuality and Intimacy

Burn survivors may have physical problems or limitations that affect sexuality and intimacy. Here are some things you can do:

- **Keep skin moisturized to lessen friction, blisters, or skin tears.** You may need to try different positioning during sexual activity or different forms of intimacy. After a burn injury, newly healed skin may be fragile and prone to shearing.
- **Good communication with your partner is important.** Newly healed areas may be sensitive to touch. Areas with skin grafts may have little or no feeling. Areas with nerve damage may be more sensitive or may have no feeling. Communicating to your partner about your trigger points may be helpful. It is important to know that it may take time through trial and error to find what works for you and your partner.

The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <http://www.msktc.org/burn/model-system-centers> for more information).



- **Consider trying different positioning or different forms of intimacy.** Contractures or tightness that limits movement, amputations, surgeries, and general weakness may have a big impact on positioning during sexual activities.
- **Talk to your doctor if you have lower energy or decreased desire.** Most survivors of a severe burn injury have lower energy levels that may improve over time. Other issues that may affect stamina include changes in hormone levels, such as decreased testosterone or increased estrogen. Medicines – particularly those that treat depression and high blood pressure – can affect desire and stamina. They can also affect a man's ability to get or keep an erection or to ejaculate. Do not stop taking any medicines suddenly because of side effects. Talk to your doctor about other medicines. Ask about blood tests to check hormone levels and nutrition status that may help your energy level and stamina after burn injury.

Psychological Effects of Burn Injury on Sexuality and Intimacy


Burn survivors may also have issues with sexuality and intimacy that relate to their mental, social, and emotional well-being. They may have reduced sexual desire and function. This may be due to the following:

- **Body image effects:** You may have changes in body image. Some individuals describe feeling:
 - Distress, shame, embarrassment, or anger. You may also feel inferior.
 - Abnormal.
 - Anxious about hiding or showing burns.
- **Emotional effects:** You may have changes in your emotional health. These may include changes in your ability to accept changes and manage your feelings as you work through challenges. These feelings may cause:
 - Fatigue.
 - Lack of sleep or changes in sleep patterns.
 - Stress and fear about having sex.
 - Mood changes.
 - Anxiety, depression, and sadness.
 - Feelings of sadness about the changes in your body.
 - Use of drugs or alcohol.
 - Boredom with your sexual routine.
- **Role change effects:** You may have changes in roles and relationships after a burn injury that can lead to:
 - A sense of not being needed, or not having control over your life.
 - Feeling like you have lost your place in your home, school, or workplace.
 - Low self-esteem due to relying on others to get through the day or due to the difficulty to fill other roles, such as being a mother, a father, a husband, a wife, or a breadwinner.
 - Ending a romantic relationship or a marriage in separation or divorce.




How can you adjust to the changes in your body after burn injury? How can you help your partner adjust to these changes?

- **You can show your interest in sex in many ways.** You may want to have your partner sit or lie next to you. Just touching and holding your partner can be fulfilling. It's important to remember that you can touch, cuddle, and caress someone without needing to have an orgasm. Sometimes just feeling loved and secure without feeling the need to perform can be enough. Over time, you will become more confident and feel more comfortable with yourself and your partner.
- **Take time to learn what you and your partner find enjoyable and exciting.** The kinds of things you did before your burn injury may not work for you now. Try to be creative and open to finding new ways to explore sexual satisfaction.
- **Be playful.** Using humor when appropriate and having fun can make your romantic relationship more interesting and enjoyable for you and your partner.



- **Keep the lines of communication open.**

- **Think about how you feel and what you want to communicate.** This will require self-awareness and self-exploration to understand what you want or need sexually after your injury.
- **Talk about your needs and wants together.** Talking about sex can be hard. Talk about things in a way that makes you both feel comfortable and secure. It might be helpful to write about your needs. The goal is to work through issues and concerns together. Your health care provider may help you resolve physical problems and concerns. Don't be afraid to ask for help.
- **Take time to listen to each other.** Good communication takes work. Listen and be open to what your partner says. Listening to each other without judgment can help solve problems in ways that satisfy both of you. Pay attention to body language. Don't assume that something will be uncomfortable; ask.
- **Be flexible.** It may take time for you to become comfortable with each other again. You may have a few setbacks. For example, you may have issues with skin sensitivity, skin breakdown, or your energy level. You and your partner will learn to manage these issues as you continue to communicate, listen, and remain flexible.



How do you keep your romance alive if your partner is your caregiver?

Be patient. Try to keep the caregiver role separate from that of a lover. This will make it easier for you and your partner to appreciate each other when you want to be intimate and allow you to keep the passion in your relationship. Here are some other suggestions:

- **Be as self-sufficient as you can.** Do as much of your self-care and other daily living activities as possible. This will help to lessen the care you need from your partner. It will also help to bring your relationship closer to how it was before your burn injury.
- **Set aside time to have fun and go on a date where you can enjoy romance without caregiving.** Keeping these roles separate will help you to avoid confusion and mix-up of the different roles.
- **Show your appreciation for each other.** Burn care can be stressful for both the patient and the caregiver.
- **If you can, hire a caregiver.** Ask a family member or friend to help with some of the daily caregiving duties. If you don't have help, see if you qualify for intervention programs that can assist you. Contact your state vocational rehabilitation agency or your church. If you are a veteran, contact the Department of Veterans Affairs.

If you are single:

- You can still have a sex life.
- If you start a new relationship, it may be hard to tell a new partner about your burn scars, and that's okay.
- You may feel anxious about hidden scars being discovered. Take your time and when you are ready, have an open and honest talk with your partner.
- Fear of rejection in a romantic relationship may affect your self-esteem. It is important to have friends who support you and listen to you.

Whether you are single or currently in a relationship—Remember the following:

- Focus on your own needs, wishes, and desires.
- Explore and communicate what gives you and your partner pleasure.
- Your sexuality is about how you feel, and your feelings may change.
- You can change your mind about how you feel, find new things that are enjoyable, and relate to your partner in new ways.
- You can aim to feel good about who you are and how you choose to experience your sexuality and intimacy.
- It's common for sex to not be on your or your partner's mind, especially in the early stages of your recovery. Some people do not want to have sex or need to have sex and that's okay too.



Where can you go with questions?

Talk to your health care provider about how the burn injury may have changed your body, your energy level, or your sense of touch. This factsheet is a starting point to help you and your partner understand that it is normal to worry about how your body may have changed after a burn injury. Each person is unique; some of your issues may be unique, too.



- Doctors who specialize in sexuality and physical and emotional trauma can help with your fears or problems. They can also give you correct information and talk openly with you. They can also make sure that you get the therapy, treatment, and information you need to help with any concerns you have about sexuality and intimacy. Your local burn center doctors can help you locate or refer you to the specialists.
- Your occupational therapist or physical therapist can be helpful as well. Occupational therapists can give you tips for how to position your body during sexual activity without causing harm or injury. Reach out to a counselor, psychologist, social worker, or sex therapist. They help people and couples identify and work through problems. They can also help you find resources.

Helpful Resources

- Your local burn center may have aftercare resources as well as events and programs to support you during recovery. For example, they may have peer support volunteers, burn survivor support groups, or survivor retreats.
- The Phoenix Society for Burn Survivors (<https://www.phoenix-society.org/>) has many resources in various formats. These include weekly chat sessions, online learning programs, workshops, and an Annual World Burn Congress.
- Factsheets and multimedia resources from the Model Systems Knowledge Translation Center on various topics related to burn injury such as *Understanding and Improving Body Image after Burn Injury*, *Social Interactions after Burn Injury*, and *Sexuality and Intimacy after Burn Injury*. Learn more at: <https://msktc.org/burn>.
- Find a provider who is familiar with sexuality topics and sexual concerns on the following websites:
 - Society for Sex Therapy and Research (STAR)—STAR is a community of professionals who have clinical and research interests in human sexual concerns. Use the link www.sstarnet.org/find-a-therapist.
 - American Association of Sexuality Educators, Counselors and Therapists (AASECT)—Search for a sexuality counselor or sex therapist in your area. Use the link www.aasect.org/referral-directory.
- Disability rights and activities organizations offer resources for sexuality for specific disabilities.



References

Alexander, M., Hicks, T., Aisen, M., & Klebine, P (2015). *Sexuality and sexual functioning after spinal cord injury*. Washington, D.C.: Model Systems Knowledge Translation Center.

Rimmer, R., & Rutter, C. (2015). Intimacy and sexuality after burn injury. *Phoenix Burn Support Magazine*, 2, 14–15.

Authorship

Sexuality and Intimacy After Burn Injury was developed by Radha Holavanahalli, PhD; Karen Kowalske, MD (North Texas Burn Rehabilitation Model System; UT Southwestern Medical Center, Dallas, TX); Ruth Rimmer, PhD, CLCP (Arizona Burn Center College of Medicine-Phoenix), and Cindy Rutter, BSN, MFTI, (Burn Survivor), in collaboration with the Model Systems Knowledge Translation Center.

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Model Systems.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this factsheet were developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Copyright © 2021 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Sleep Problems after Burn Injury

October 2020

www.msktc.org/burn/factsheets

BURN Factsheet

This fact sheet explains the causes of sleep problems after burn injury and provides treatment options.

Introduction

Sleep problems occur in more than 50% of people who have had severe burn injuries. Insomnia is the most common type of sleep problem. Insomnia can take many forms, including:

- Difficulty falling asleep.
- Difficulty staying asleep.
- Poor quality sleep.
- Waking too early.
- Nightmares.

Sleep problems are especially common right after burn injury and during the healing and recovery stages. Insomnia can come and go over the years and may require different solutions at different times. Everyone has a different experience after burn injury, and some of the following information may not apply to you.

Causes of Sleep Problems after Burn Injury

Many factors can disturb sleep after burn injury. Some may continue to affect you long after leaving the hospital and healing.

- Anxiety and post-traumatic stress disorder (PTSD), including fear of falling asleep to avoid nightmares or to remain on the alert against threat.
- Depression.
- Constantly thinking about the burn event.
- Pain.
- Itching.
- Many medications prescribed to treat any of the above problems can also disturb sleep.
- Burn injury can change hormone levels and other chemicals in the body that regulate or affect sleep.
- Sleep apnea (pauses in breathing during sleep)—symptoms include day-time sleepiness, snoring, and agitation.
- Contractures caused by scar tissue—this can limit the ability to move and get comfortable.
- Difficulty breathing, if the respiratory passages (nose and throat) were affected by the burn injury.
- See your health care providers regularly and follow through with treatment. Success with getting back to work requires active participation on your part.

The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <http://www.msktc.org/burn/model-system-centers> for more information).



Model Systems
Knowledge Translation
Center

Poor Sleep Can Cause Serious Problems

It is important to get treatment for poor sleep because it can be harmful in a number of ways. It can be extremely distressing and debilitating, and actually interfere with your recovery from the burn injury. Poor sleep can:

- Make pain worse
- Slow down wound healing.
- Cause restlessness, irritability, and changes in behavior.
- Cause problems during the day such as:
 - Low mood or depression.
 - Trouble handling stress.
 - Lack of energy.
 - Difficulty concentrating.
 - Increased risk of accidents.

Role of Melatonin

Melatonin is a natural hormone made by your body that is released into your blood stream to trigger sleepiness. A decrease in light signals the body to release this hormone. The light from electronic screens (blue light) such as that from your TV, computer, smart phone, or electronic books can interfere with the production of melatonin. These devices should be turned off at least one hour before bedtime. Melatonin can also be taken by mouth to promote sleep. Talk to your doctor about this treatment option.

Treatment Options

There are many different approaches to solving sleep problems. The choice of treatment depends on the cause, type, and severity of the problem as well as your stage of recovery from burn injury. Your doctor or medical team may talk to you about your past sleep habits and other factors that affect sleep in order to better understand your sleep problems. If necessary, your doctor may refer you to a sleep specialist.

Treatments That Do Not Involve Medications (also called behavioral or nonpharmacological treatments)

Good Sleep Hygiene

- Sleep hygiene is the practice of following sensible guidelines for promoting regular, restful, good-quality sleep.
- Daytime naps can disrupt normal sleep patterns. Naps may be needed in early stages of recovery, but should gradually decrease as your health improves.
- If naps are needed, they should be at the same time every day and last no more than 30 minutes.

- Maintain a regular time for getting up in the morning, regardless of when you fell asleep. If you have trouble falling asleep after 15 minutes, you should get out of bed and do something else (reading, word puzzles, listening to music) until you get sleepy.
- Do not go to bed until you are tired. Establish a consistent bedtime routine to cue your brain that it is time for sleep.
- Avoid stimulant-containing drinks, food, and drugs in the late evening. Caffeine is in many soft drinks, chocolate, candy, coffee, and bakery products.
- Steer clear of stimulating activities late in the evening, such as surfing the Web, watching exciting or frightening movies, or playing video games.
- Start increasing physical activity and exercising regularly as soon as your doctor says it is okay. Exercise is especially helpful if you suffer from anxiety as well as trouble sleeping.
- Vigorous exercise should be done earlier in the day, at least 6 hours before bedtime.
- Mild exercise should be done at least four hours before bedtime.
- Don't go to bed hungry, but avoid large meals close to bedtime. If you have trouble staying asleep, a light snack an hour before bedtime may help you sleep through the night.
- Alcohol can cause you to sleep poorly. Also, alcohol can be dangerous if you are already taking medications that make you drowsy. Ask your doctor if it is safe for you to drink alcohol.
- Avoid smoking or using other forms of nicotine close to bedtime. Nicotine is a stimulant.

Stimulus Control

If trouble falling asleep goes on for a long period of time, sometimes people develop a “habit” of thinking they won’t fall asleep, and these thoughts keep them awake. Stimulus control can help “re-program” you to associate the bedroom and bedtime with only sleep-promoting (calm and pleasant) activities rather than failure to fall asleep.

Guidelines for stimulus control:

- Go to bed only when sleepy.
- Set a regular wake-up time, no matter what time you actually fall asleep the night before.
- Get out of bed whenever you are awake for longer than 15-20 minutes.
- Avoid reading, watching TV, eating, or worrying in the bed and bedroom.
- Do not nap during the day.

Relaxation Training

Relaxation techniques reduce anxiety and tension at bedtime to help you fall asleep. They can also be used to fall back asleep if you awake in the night. There are several techniques.

- Progressive muscle relaxation
- Meditation training



Model Systems
Knowledge Translation
Center

- Imagery training
- Biofeedback
- Hypnosis
- Yoga

For any of these methods, a practitioner will teach you the formal steps or exercises involved. These methods are most successful when practiced regularly at home. Apps can also be downloaded onto a smart phone or tablet that promote sleep by using relaxing sounds, relaxing music, or even sleep casts or sleep stories. Many of these apps are free.

Cognitive Behavioral Therapy (CBT)

Cognitive behavioral therapy for insomnia (CBTi) teaches sleep hygiene, employs stimulus control techniques, and trains in relaxation methods/practices. In addition, sometimes people with sleep problems develop thoughts and beliefs about sleep that keep them from falling asleep. Examples of such thoughts include “I will never be able to fall asleep,” or “I will not be able to function tomorrow since I cannot fall asleep,” or “I cannot sleep well without alcohol.” A cognitive behavioral therapist can work with you to address and eliminate the thoughts that may be keeping you from being able to fall asleep.

Light Therapy

Our body’s chemicals and hormones vary in a natural 24-hour cycle that promotes sleep or wakefulness at certain times. If this natural cycle (called “circadian rhythm”) gets disturbed for any reason, sleep problems can occur. Light therapy uses exposure to daylight (or “light boxes” that mimic daylight) to “reset” the circadian rhythms for sleeping and waking.

- Different ways to do this are using light boxes in the morning, taking daytime walks outside, or using light machines called “dawn stimulators” that mimic a gradual sunrise.
- Light therapy is often used along with sleep hygiene.

Medication (Pharmacologic) Treatment

There are effective medications that can help you sleep better. These may include sleep aids, antidepressants and/or anxiety medications. They can be used alone or in addition to one of the above approaches to improve sleep. As with any medications, it is extremely important to take medications for sleep only as your doctor has prescribed and discussed with you. This includes over-the-counter sleep medications.



Model Systems
Knowledge Translation
Center

Bibliography

Mayes T, Gottschlich MM, Khoury J, McCall J, Simakajornboon N, Kagan R. A pilot review of the long-term impact of burn injury on sleep architecture in children. *J Burn Care Res.* Jan-Feb 2013;34(1):e15-21. Lee AF, Ryan CM, Schneider JC, et al. Quantifying risk factors for long-term sleep problems after burn injury in young adults. *J Burn Care Res.* 2017 Mar/Apr;38(2). E510-e520.

Jaffe SE and Patterson DR. Treating Sleep Problems in Patients with Burn Injuries: Practical Considerations, *Journal of Burn Care & Rehabilitation.* 2004 May-Jun; 25(3):294-305

Harorani M, Davodabady F, Masmouei B, Barati N. The effect of progressive muscle relaxation on anxiety and sleep quality in burn patients: A randomized clinical trial. *Burns* 2020 (46):1107-1113.

Authorship

Sleep Problems after Burn Injury was developed by Shelley A. Wiechman, Ph.D., in collaboration with the Model Systems Knowledge Translation Center.

Factsheet Update

Sleep Problems after Burn Injury was reviewed and updated by Shelley A. Wiechman, Ph.D., in collaboration with the Model Systems Knowledge Translation Center.

Source: The content in this factsheet is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Model Systems (BMS), funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR).

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. This publication was produced by the Burn Model Systems in collaboration with the University of Washington Model Systems Knowledge Translation Center with funding from the National Institute on Disability and Rehabilitation Research in the U.S. Department of Education, grant no. H133A060070. It was updated under the American Institutes for Research Model Systems Knowledge Translation Center, with funding from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Copyright © 2020 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Social Interaction After Burn Injury

June 2019

www.msktc.org/burn/factsheets

BURN Factsheet

This factsheet explains challenges with social interactions after burn injury. It describes some strategies you can use when meeting new people or going into new social, work, school, or public situation.

Some burn injury survivors have changes in their appearance. They may worry about how people will react to them when they leave the hospital and go out in public.

Understanding challenges with social interactions

Some of the social challenges that burn survivors face after they leave the safe zone of the hospital and return to their communities include

- Stares, startled glances, or double-takes;
- Nosy questions or comments about their injury; and
- Bullying and teasing.

If your burns show, everyday activities like going shopping or taking a bus or the subway may involve being stared at and having to deal with people's curiosity. If your burns are hidden, you may worry about how people will react when they are exposed, such as when you take off your shirt at the beach.

The way people react can make it hard to feel confident during social interactions. People may react verbally, with words. People may also react nonverbally, through body language or gestures. Some burn survivors aren't bothered by the reactions of others. But you may find it helpful to learn skills to face these challenges.

Do you sometimes feel uncomfortable in social situations?

Some burn survivors may avoid social situations because they think that people may react to their appearance. Other burn survivors may

- Feel nervous or on edge when they meet new people or when they're with strangers;
- Feel isolated and alone; or
- Feel emotional distress, including depression.

Finding help to improve social interactions

If you feel nervous or on edge in social situations, you can learn skills to help you gain control, comfort, and confidence. Some resources that may help include:

- Members of your burn team can talk with you and provide resources to help you.

The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <http://www.msktc.org/burn/model-system-centers> for more information).



Model Systems
Knowledge Translation
Center

- The Phoenix Society for Burn Survivors has a peer support program called Survivors Offering Assistance in Recovery (SOAR) for burn survivors and their loved ones. This one-on-one program connects people with new burn injuries to other survivors and family members who can share their experiences and offer support.
- The Phoenix Society also offers an online learning program called Beyond Surviving: Tools for Thriving After Burn Injury. This program teaches strategies to help you gain control, act and feel comfortable, and confident.
- The Phoenix Society also offers a weekly peer support chat. This live chat is held on Wednesday nights from 9:00 p.m. to 10:30 p.m. (EST). It covers common concerns and questions facing burn survivors and their loved ones. See <https://www.phoenix-society.org/?modal=what-we-do> for more information.

It's okay to feel nervous when interacting with people after a burn injury. Here are some strategies you can use when meeting new people or going into new social, work, school, or public situations. In all social interactions, it's helpful to act positive and use confident body language.

The “STEPS” strategy may help you feel more confident when facing old or new social, work, and school situations. It was developed by Barbara Kammerer Quayle, a burn survivor.

- **Self-talk**—This is what you say to yourself and believe. Examples include “I love and accept myself the way I am and the way I am not,” “I meet people easily and feel comfortable with them,” and “I can do it!”
- **Tone of voice**—Use a friendly, warm, and enthusiastic tone of voice.
- **Eye contact**—Look people in the eye, even if just for a few seconds.
- **Posture**—Have a confident posture. Keep your head up, your rib cage lifted, and your shoulders back.
- **Smile**—A smile makes you look confident and approachable.

Ways you can respond when others react to you

If someone stares at you

People who stare are often just curious because they haven't seen someone with a burn injury. They usually don't mean to be rude. If you look back and smile and say, “Hi” or “Hi, how are you doing?” the staring usually stops. Always use your STEPS to look confident and comfortable.

If someone asks what happened to your burn injury

When strangers ask about your burn injury, remember that you are in control. You can offer as much or as little information as you want. You don't owe strangers the whole story of your burn injury experience.

Rehearse your responses. This requires writing and then saying three short sentences about your injury, smiling at the person asking, and walking on. For example:

- I was burned in a house fire.
- I am doing better now, or I am still recovering.
- Thanks for your concern, or thanks for asking.

Write and rehearse three sentences that feel just right for you and use this tool when people ask questions.

Respond in a positive way.

Change the subject if you don't want to continue to talk about your burn.

Use humor when appropriate to lighten the moment.

You can find more information about the Rehearse Your Responses strategy at <https://www.phoenix-society.org>. Look for: Find Resources/Beyond Surviving Tools for Thriving.

If someone teases you:

- Stand up straight and hold your head high.
- Count to 10 slowly and stay calm.
- Shrug your shoulders, act bored, smile, and walk away.
- Say to the teaser, "I'm wondering why you would say something like that?" or "you must be pretty insecure to pick on other people." Shrug your shoulders and walk away.

Social Skills Training

Social interaction skills training programs may also help you

- Get ready for social situations after you leave the hospital
- Understand what goes on in social interactions
- Practice effective ways to manage social interactions

Two well-known social interaction skills training programs are described below.

- Beyond Surviving: Tools for Thriving After a Burn Injury," (<https://www.phoenix-society.org>) which is available through the Phoenix Society and described in this factsheet
- The **3-2-1-GO! strategy**—This program was developed by James Partridge (<https://www.changingfaces.org.uk>). It includes
 - Three things to do if someone stares at you
 - Two things to say if someone asks what happened
 - One thing to think if someone turns away

Resources

The Phoenix Society, Inc.
525 Ottawa Ave NW, Front
Grand Rapids, MI 49503

(800)888-2876 or (616) 458-2773

info@phoenix-society.org

Changing Faces The Squire Centre
33-37 University Street London, WC1E 6JN
<https://www.changingfaces.org.uk>

References

Blakeney, P., Partridge, J., & Rumsey, N. (2007). Community integration. *Journal of Burn Care and Research*, 28(4), 598–601.

Blakeney, P., Thomas, C., Holzer, C., 3rd, et al. (2005). Efficacy of a short-term, intensive social skills training program for burned adolescents. *Journal of Burn Care & Rehabilitation*, 26(6), 546–555.

Kammerer Quayle, B. (2006). Behavioral skills and image enhancement training for burn survivors: Essential interventions for improving quality of life and community integration. In R. Snood & B. Achauer, (Eds.), *Achauer and Snood's burn surgery: Reconstruction and rehabilitation*. Elsevier, Inc.

Kammerer Quayle, B. (2015). Program that has helped thousands began with one survivor's need to get back to living. *Burn Support Magazine*, 2.

Authorship

Social Interaction After Burn Injury was developed by Radha Holavanahalli, PhD and Barbara Kammerer Quayle, MS in collaboration with the Model Systems Knowledge Translation Center.

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Model Systems.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. This publication was produced by the Burn Model Systems in collaboration with the University of Washington Model Systems Knowledge Translation Center with funding from the National Institute on Disability and Rehabilitation Research in the U.S. Department of Education, grant no. H133A060070. It was updated under the American Institutes for Research Model Systems Knowledge Translation Center, with funding from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this fact sheet do not necessarily represent the policy of the U.S. Department of Education or the U.S. Department of Health and Human Services, and you should not assume endorsement by the federal government.

Copyright © 2019 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Sun Protection After a Burn Injury


May 2023

<https://msktc.org/burn/factsheets>

BURN Factsheet

This factsheet explains the importance of sun protection after burn injury. It describes how sun exposure affects your healed skin and grafts and ways to limit sun exposure. It also provides resources to learn more about protecting your skin.

The sun has many beneficial properties, but the sun emits three types of ultraviolet (UV) light that can harm skin and has been linked to skin cancers (basal and squamous carcinoma, melanoma) and vision problems (cataracts and macular degeneration). Healed burns, donor sites and skin grafts are more sensitive to sunlight.



Your Burn Injury and Sun Exposure

Burn skin sensitivity. Healed burns or skin grafts may be extremely sensitive to sunlight and may sunburn more severely even after short periods of time in the sun compared to before your injury. Sun sensitivity after a burn injury may last for a year or more. In addition, some medications can cause you to be more sensitive to the sun (e.g., anti-itch medicines, like cetirizine, and non-steroidal anti-inflammatory medications, like ibuprofen).

Pigmentation. The color of our skin is related to the amount of melanin that each of us has in our skin. When someone has a second degree or deeper burn injury, the pigment of the skin is initially lost because melanin is located in the epidermis (the outer most layer of skin). With healing, the pigment may return, but this process is unpredictable. Often, newly healed skin appears pink and unpigmented regardless of the original skin tone. As the scar matures, the skin may regain normal pigment, regain more pigment than it had before the injury, or stay unpigmented. The deeper the burn wound, the slower the re-pigmentation process. There is no way to predict if there will be a color difference after the healed skin has matured. Some people will have lighter (called hypo-pigmented) or darker (called hyper-pigmented) skin than they had in that area before the injury.

The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <https://msktc.org/burn/model-system-centers> for more information).

Because re-pigmentation can be affected by UV light (sunlight), suntans in newly healed skin may not fade when the rest of the tan goes away. For a burn that needs a skin graft to heal, the grafted area may become darker than the surrounding uninjured skin. The reason for this is not clear, but sun protection is encouraged to prevent pigment changes that may result in permanent skin tone changes within the burn wound for at least one year after a burn injury.



People with darker skin. Even if you have never had issues with sun exposure or sunburn in the past, your areas that were burned or skin grafted are now susceptible to sunburn and pigmentation changes. Sun exposure will not necessarily increase pigmentation in areas of your scar that have less pigment than your surrounding skin.



Model Systems
Knowledge Translation
Center

How Can You Protect Your Skin From Exposure to the Sun?

Returning to your normal activities (e.g., walking or playing outside, hiking, swimming, biking, gardening) after a burn injury is important and strongly encouraged. However, if you plan to be outside for long periods of time, using a combination of methods of sun protection is the best way to protect your skin. Dermatologists (skin doctors) recommend using a combination of sun avoidance, protective clothing, and sunscreen/sunblock to limit sun/UV light exposure and to combat sun damage.



Sun Avoidance

It is nearly impossible to stay out of the sun and be active. But you can choose when you are outside. The sun is strongest from 10 a.m. to 4 p.m. or when your shadow is shorter than your height. Don't let clouds fool you. Up to 80% of harmful UV sunrays can penetrate clouds. Also, keep in mind that UV rays reflect off sand, water, snow, and ice and can cause additional UV sunray exposure. Certain types of UV light are stronger at higher altitudes. Thus, limiting peak sun exposure and being aware of your environment are keys to preventing sun damage to your skin. Seek shade when you are outside.

What about tanning beds? Tanning beds should be avoided, as they emit the same amount (if not more) cancer-causing UV light. And they will cause pigmentation that might not fade in healing scars.

Protective Clothing

Protective clothing is a good way to protect yourself and your healed skin from the sun. It is important to know that not all fabrics and clothing are protective. When choosing clothing to provide protection from harmful UV rays, consider the following:

- The tighter the weave of the fabric, the better the protection. For example, fabrics such as denim and wool have a tighter weave and provide better protection than lighter fabrics like linen.
- Loose clothing tends to provide better protection than tight or stretched clothes.
- Wet fabrics provide less protection than dry fabrics.
- Synthetic and semi-synthetic materials tend to provide better protection than bleached or refined cottons.
- Dark clothing provides better protection.
- Specialized, sun-protective clothing made from fabric that has been embedded with chemicals that deter or absorb harmful UV rays is increasingly available at sporting stores and online. An Ultraviolet Protective Factor (UPF) rating system that is listed on the tag describes the degree of protection the clothing provides against UV light. Specialized clothing with ratings of UPF 30 or higher is recommended.

Other Protective Wearable Options

Other types of protective options include wearing sunglasses and wide-brim hats. Recommendations include:

- Sunglasses with 99–100% UV protection that completely cover the eyes and eyelids.
- Hats should have at least a 3-inch brim all the way around to protect the neck, ears, and face. Baseball caps do NOT provide adequate protection for facial and neck burns.
- Hats with a neck flap option have better coverage of the neck.



Sunscreens

Sunscreens are a third line of defense. Keep in mind that sunscreens should be used **TOGETHER** with protective clothing and sun avoidance. A sunscreen's ability to protect you against dangerous sunrays is measured by a number called SPF or Sunburn Protection Factor. SPF refers to how long it takes for skin with sunscreen to burn compared to skin without sunscreen. Generally, the higher the SPF number, the greater protection against sunburn.



Sunscreens come in multiple forms: gels, sprays, creams, and sticks. Gels tend to work best on hairy areas, while creams work best on the face and dry skin. Sticks work well around the eyes and lips. FDA regulations and standardizations do not apply to spray sunscreens; it is important that the spray covers all exposed skin, is not inhaled or used near a heat source, while smoking or near an open flame. Some sunscreen products contain alcohol, which can be drying to recently healed skin or grafted areas. Sunscreens are also available within face creams and make-up (e.g., foundations). Although convenient, they must be applied frequently to achieve the most sun protection.

Some insect repellents also contain sunscreen. The American Academy of Dermatology recommends that insect repellent be applied separately from sunscreen, as the repellent should be applied minimally and sparingly, while sunscreens require generous and frequent applications.

What is the difference between a sunblock and sunscreen? Sunscreens are more common, and they filter or screen the sun's ultraviolet rays. They keep most rays out but do let some in. A sunblock reflects the sun's rays and is considered a physical barrier. Most sunblocks use titanium dioxide or zinc oxide, which can make them thicker and somewhat opaque (not transparent) when applied to the skin. Many products commonly referred to as sunscreens combine agents.

Can I use a suntan lotion instead? No. Often the SPF levels in a suntan lotion are only 4 or 5, which is **not** enough to protect your skin from the sun.

Many studies have shown that people apply sunscreen incorrectly, resulting in inadequate protection. The following tips regarding sunscreen use will help you avoid a sunburn and sun damage:

- Apply sunscreen every day, all year, even on overcast or cloudy days.
- Use a sunscreen with SPF 30 or greater.
- Use a broad-spectrum sunscreen to protect you from both UV-A and UV-B light.
- Apply or reapply every 2 hours when outside.
- Use water-resistant sunscreens.
- Generally, one handful of sunscreen lotion covers all uncovered areas of your skin.
- Keep newborns and babies younger than 6 months out of direct sunlight. Use of sunscreen on babies younger than 6 months is not recommended as they are at greater risk for sunscreen side effects, such as a rash.

*Experts do not recommend one type over the other as long as the product **protects your skin from both UV-A and UV-B sun rays, has an SPF of 30 or higher, is water resistant, and is applied according to the directions on the container.***



- For babies older than 6 months, The American Academy of Pediatrics recommends using sunscreen on infants for small areas such as the face and back of the hands where protection from clothes is inadequate. Generally, products with zinc oxide or titanium dioxide are less irritating.
- Apply sunscreen 30 minutes before going outside to allow the sunscreen to absorb into your skin.
- Reapply sunscreen every 2 hours and immediately after swimming/sweating heavily/drying yourself with a towel. This also applies to the use of water-resistant sunscreens.
- Cover your lips with a sunscreen-containing lip balm of SPF 30 or greater and reapply frequently. Avoid petroleum jelly-based products that do not include sun protection.
- Don't forget about your ears, feet, back of your hands, neck, and bald spots when applying sunscreen.
- Sunscreen designed for the face usually does not clog pores and cause pimples and may be better than other sunscreens. Moisturizers with sunscreen are available and provide both hydration and sun protection; choose moisturizers with SPF 30 or higher.



Frequently Asked Questions About Sunscreen Use

- *Will it ever be safe for me to wear short sleeve shirts and shorts again?* We understand that in hot weather long sleeves and pants can be uncomfortable. Short sleeves and bathing suits are OK if you are careful about using sunscreen and limiting how much time you spend in the sun.
- *What should I do if I have a reaction (like a skin rash) to sunscreen?* Stop using the product immediately and try another one with different ingredients. Talk to your doctor or healthcare provider if the reaction persists.
- *How long is my bottle of sunscreen good for?* Based on FDA reports, sunscreens are typically good for 3 years. Many sunscreens will have an expiration date on the bottle. If there is no date, write the purchase date on the bottle and discard after 3 years. If the consistency or texture of the sunscreen changes, discard the bottle.
- *What is a PABA-free sunscreen?* PABA is the abbreviation for para-aminobenzoic acid and related chemicals. PABA-containing sunscreens was an UV-B absorbing type of sunscreen used in the early 1940s. Since then, PABA has been removed from most sunscreens due to reports of allergies and reactions. PABA-free sunscreens do not contain PABA or any related chemicals.
- *Does a higher SPF sunscreen work better than a lower SPF?* No sunscreen blocks 100% of UV sun rays. SPF 30 blocks 97% of UV light when used appropriately. The increased SPF does not imply that you can use a smaller amount or apply it less frequently. All sunscreens should be applied generously every two hours or as directed on the container, no matter the SPF number.
- *Can sunscreens cause cancer?* Recent studies have shown no link between sunscreen use and melanoma (a form of skin cancer).
- *Are sunscreens toxic?* Sunscreens undergo extensive testing to ensure that they do not cause harm to humans. The safety of the chemicals used is based on studies with humans but not specifically with people.
- *Do sunscreens cause Vitamin D deficiency?* Sunscreens theoretically prevent Vitamin D formation because they block the UV-B light rays that are needed to form Vitamin D. However, no studies have shown any physical consequence or difference in Vitamin D levels of people who use sunscreen when compared to people who do not use sunscreen. We encourage a healthy diet that is rich in Vitamin D (fatty fish, cheese, egg yolk, fortified milk and yogurt, beef liver).



- *Am I more susceptible to skin cancer after a burn?* There is no good information about the risk of skin cancer with healed burn wounds.
- *When should I be concerned about skin cancer?* You should “check your birthday suit on your birthday” for skin changes or itchy or bleeding skin. If you have a burn wound that does not heal or breaks open often you should see your doctor or health care provider.

Other Resources

- Phoenix Society: <https://www.phoenix-society.org/resources/sun-protection-after-burns>
- American Cancer Society: <https://www.cancer.org/cancer/skin-cancer/prevention-and-early-detection.html>
- Skin Cancer Foundation: <http://www.skincancer.org/prevention/sun-protection/clothing/clothing-our-first-line-of-defense>
- FDA Factsheet: *Tips to Stay Safe in the Sun: From Sunscreen to Sunglasses*: <http://www.fda.gov/ForConsumers/ConsumerUpdates/ucm049090.htm>
- American Academy of Dermatology: Sunscreen FAQs: <https://www.aad.org/media/stats-sunscreen>
- U.S. Environmental Protection Agency: Sunscreen—The Burning Facts: <https://www.epa.gov/sites/production/files/documents/sunscreen.pdf>



Authorship

Sun Protection After a Burn Injury was originally developed by Gretchen J. Carrougner MN, RN; Nicole S. Gibran, MD; Maria Caceres, BSN, RN; and Cathie Cannon RN in collaboration with the Model Systems Knowledge Translation Center (MSKTC). It was reviewed and updated by Gretchen J. Carrougner MN, RN in collaboration with the MSKTC.

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems. Because of limited research on sun exposure following a burn injury, this factsheet was developed based on studies and information for the general population.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this factsheet were developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082) and were updated under an NIDILRR grant (90DPKT0009). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, HHS, and you should not assume endorsement by the Federal Government.

Recommended Citation: Carrougner, G. J., Gibran, N. S., Caceres, M., & Cannon, C. (2023). *Sun protection after a burn injury*. Model Systems Knowledge Translation Center (MSKTC). <https://msktc.org/burn/factsheets/sun-protection-after-burn-injury>.

Copyright © 2023 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Temperature Sensitivity After Burn Injury

August 2023

<https://msktc.org/burn/factsheets>

BURN Factsheet

This factsheet explains the causes of temperature sensitivity, how it may affect your life, and tips to cope with it.

What Is Temperature Sensitivity?

Many people deal with temperature sensitivity after a burn injury. Temperature sensitivity is a broad term that includes several issues. Some people may have issues when they touch warm or cool objects or liquids. Others may have challenges regulating their body temperature or tolerating extremes of temperatures or air movement; the ability of your body to control its temperature is called thermoregulation.



Temperature sensitivity is more common among people who had flame burns or large burns, those whose scars are in areas that are more exposed (e.g., hands, face), or those who had more skin grafting. Desensitization is a way to re-train your nerves by slowly exposing yourself to more and more intense textures, pressures and temperatures.

Causes of Sensitivity to Warm or Hot Environments

One way your body cools itself down is to increase blood flow to the skin, which lets heat out. Grafted and scarred skin does not do this as well as uninjured skin. Another way your body cools itself is through sweating, which cools the skin. Grafted and scarred skin may not sweat like uninjured skin due to loss of sweat glands. If your skin can't perform these methods of cooling, you will be more sensitive to warmer temperatures. Conversely, uninjured skin may sweat excessively as it tries to overcome the lack of sweat from grafted or scarred skin. Some tips for cooling are listed below.

Causes of Sensitivity to Cold Environments

Cold temperatures can trigger nerve pain and cause skin to dry out and crack. It can be challenging for people living with burn injuries to be outside when it's cold or windy. These environments may even cause pain that requires accommodations like those described below.



When it's cold, our bodies decrease blood flow to the skin, which limits heat loss from our bodies. Although this may initially be impaired, grafted or scarred skin may be able to do this normally over time.

Causes of Touch Sensitivity and Numbness

Neuropathic pain, or nerve pain, commonly occurs after burn injury. Damage to and re-growth of nerve endings in your skin can cause this type of pain. Nerve endings sense pain, itch, pressure, sharpness, and temperature. They also send signals to your brain to tell you that something feels hot, cold, or painful. When nerves are injured and recovering, these signals can increase and are sometimes

The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <https://msktc.org/burn/model-system-centers> for more information).



Model Systems
Knowledge Translation
Center

incorrect. For example, your nerves may sense pain for usually non-painful stimuli like air or clothes moving across a wound, graft, or scar. This leads to more intense hot or cold feelings than you would get from uninjured skin or pain from touching something that shouldn't cause pain.

Damaged nerve endings don't always repair themselves fully. In these cases, you may feel less sensation to hot, cold, or pain, or even numbness. These signals normally let you sense when something is too hot or too cold, which triggers a reflex to pull your hand or exposed body part away. This reflex helps to prevent injury from intense temperatures. When you can't feel temperature or pain, you will not have the same reflex. This puts you at greater risk of hurting yourself on surfaces that are too hot or cold, leading to reinjury of your burn scars. You must pay special attention to high-risk situations and injured areas, for example, when using a heating pad or soaking feet in warm water.



Donor Sites and Temperature Sensitivity

A donor site, the area where skin is taken for skin grafts, should keep relatively normal blood flow and sweat production after it heals. This means that it should be able to help regulate your body temperature to the surrounding air temperature.

How Can Temperature Sensitivity Affect Your Life?

Impact of Warm and Cold Environment Temperature Sensitivity

- Becoming overheated or cold because of issues with regulating your internal body temperature.
- Overheating is dangerous and can cause heat exhaustion and heat stroke. If you start to feel a headache, nauseated, dizzy, weak, or especially thirsty get help and cool down immediately.
- Increased sweating of the uninjured skin that is trying to compensate for the skin and scar that can no longer sweat normally when hot.
- Trouble doing daily activities in hot or cold temperatures. This includes social activities.
- If you work outside, temperature sensitivity can make it hard to go back to work or to do pre-injury tasks. People often require accommodations at work, like being able to take additional breaks, cool down, or change clothes. Some people need to work in different environments (e.g., indoors during the summer or winter when they worked outdoors year-round before the injury).

Impact of Touch Sensitivity

- It can be painful and uncomfortable to touch surfaces and liquids that are warm or cool. This requires focused retraining and desensitization. Talk with your burn team about this process.
- The feeling of pain can lead to avoidance of contact and fear of physical movement. This is known as kinesiophobia.

Tips for Coping With Temperature Sensitivity

Warm or Hot Environment Sensitivity Tips

- Plan your day so that you are active during the cooler times.
- If you work outside, talk with your employer about your comfort and safety needs.
- Add a few drops of peppermint essential oil to your moisturizer.



- Wear light, synthetic, breathable fabric.
- Wear a wide brim hat out in the sun.
- Wear layers so you can easily adjust to changes in temperature.
- Stay well hydrated.
- Use wearable cooling devices such as a cooling vest, a cooling hat insert, or a neck fan.
- Move to a cool location before you begin to overheat.
- Mist yourself with cool water if you are feeling overheated.
- If you need to cool down fast, find air conditioning and/or fans and use cold packs and damp cloths on your forehead or neck and call for help.
- Work with a therapist to learn relaxation, meditation, and visualization techniques. This may, for example, help you visualize yourself in a cooler place or in a place where you feel more comfortable when you are experiencing discomfort from heat.



Cold Environment Sensitivity Tips


- Wear a base layer made of fabric like wool or synthetic material that wicks sweat away from your skin.
- Wear layers so you can adjust to changes in temperature.
- Cover any areas that are sensitive to protect them from wind and cold air. You may need gloves, a hat or a ski mask.
- Slowly adjust to the cold. For example, gradually exposing yourself to cold air daily.
- Use thicker moisturizers to act as a barrier to cold and wind. They can also help with the dryness and tightening of skin and scars.
- If you work outside, talk with your employer about your comfort and safety needs.

Touch Sensitivity Tips

- Talk to your health care provider about creating a personal pain management plan to help you cope with pain. A pain management plan may include a mix of medicine and behavioral approaches. Examples of behavioral approaches include breathing, visualization and mindfulness techniques.

Resources

- Contact your health care provider or burn team for more information and for help coping with temperature sensitivity.
- Reach out to support groups or other burn survivors for additional help coping with temperature sensitivity.
- Resources on itchy skin after burn injury: <https://msktc.org/burn-topics/itchy-skin-after-burn-injury>
- Resources on managing pain after burn injury: <https://msktc.org/burn-topics/managing-pain>
- Resources on sun protection after burn injury: <https://msktc.org/burn/factsheets/sun-protection-after-burn-injury>
- Phoenix Society for Burn Survivors: 1-800-888-2876: <https://www.phoenix-society.org/>



References

Oh, J., Madison, C., Flott, G., Brownson, E. G., Sibbett, S., Seek, C., Carrougher, G. J., Ryan, C. M., Kowalske, K., Gibran, N. S., & Stewart, T. (2021). Temperature sensitivity after burn injury: A burn model system national database hot topic. *Journal of Burn Care & Research*, 42(6), 1110–1119.



Model Systems
Knowledge Translation
Center

- Davis, S. L., Shibasaki, M., Low, D. A., Cui, J., Keller, D. M., Purdue, G. F., Hunt, J. L., Arnoldo, B. D., Kowalske, K. J., & Crandall, C. G. (2007). Impaired cutaneous vasodilation and sweating in grafted skin during whole-body heating. *Journal of Burn Care & Research*, 28(3), 427–434.
- Ramanlal, R., & Gupta, V. 2022. *Physiology, Vasodilation*. StatPearls Publishing.
<https://www.ncbi.nlm.nih.gov/books/NBK557562/>
- Davis, S. L., Shibasaki, M., Low, D. A., Cui, J., Keller, D. M., Purdue, G. F., Hunt, J. L., Arnoldo, B. D., Kowalske, K. J., & Crandall, C. G. (2008). Cutaneous vasoconstriction during whole-body and local cooling in grafted skin five to nine months postsurgery. *Journal of Burn Care & Research*, 29(1), 36–41.
- Cramer, M., Morales, G., Huang, M. U., & Crandall, C. G. (2019). No thermoregulatory impairment in skin graft donor sites during exercise-heat stress. *Medicine & Science in Sports & Exercise*, 51(5), 868–873.
- Williams, F. (2021). *Ask the experts: extreme cold sensation*. Phoenix Society for Burn Survivors.
<https://www.phoenix-society.org/resources/ask-the-experts-extreme-cold-sensation>
- O'Brien, C., Castellani, J. W., & Sawka, M. N. (2011). Thermal face protection delays finger cooling and improves thermal comfort during cold air exposure. *European Journal of Applied Physiology*, 111(12), 3097–3105.
- Gobel H, Schmidt G, Dworschak M, Stolze H, Heuss D. (1995). Essential plant oils and headache mechanisms. *Phytomedicine*, 2(2), 93-102.

Authorship

Temperature Sensitivity After Burn Injury was developed by Caitlin Orton, MPH, Jamie Oh, MD, Grace A. Flott, Craig Crandall, PhD, and Karen Kowalske, MD, in collaboration with the Model Systems Knowledge Translation Center.

Source: The content in this factsheet is based on research and/or professional consensus. This content has been reviewed and approved by experts from the Burn Model System (BMS) centers, funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR). The content of this factsheet has been reviewed by burn survivors and/or their families.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider about specific medical concerns or treatment. The contents of this factsheet were developed under grants from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant numbers 90DPKT0009 and 90DPBU0005). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the federal government.

Recommended Citation: Orton, C., Oh, J., Flott, G. A., Crandall, C., & Kowalske, K. (2023). *Temperature sensitivity after burn injury*. Model Systems Knowledge Translation Center (MSKTC).
<https://msktc.org/burn/factsheets/temperature-sensitivity-after-burn-injury>

Copyright © 2023 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Understanding a Burn Injury

September 2023

<https://msktc.org/burn/factsheets>

BURN Factsheet

This factsheet offers important details about burn injuries to burn patients who are admitted to hospitals. This content may also be helpful to the family members and friends of burn patients.


PHASE I: Burn Evaluation & Early Care

What Is a Burn Injury?

A **burn injury** is damage to the layers of skin caused by exposure to the one of the below methods. Deeper burn injuries may damage tissue (fat and muscle), or even bone.

- **Flame:** contact with fire or flames
- **Flash:** resulting from the heat of an explosive blast
- **Scald:** contact with hot liquids or hot steam
- **Grease:** contact with hot grease
- **Contact:** prolonged contact with something hot
- **Electrical:** electricity passing through the body and heating the skin and underlying tissue
- **Chemical:** contact with chemicals, such as acid or alkalis

What Are the Different Degrees of Burn Injury?



The Burn Model System Program is sponsored by the National Institute on Disability, Independent Living, and Rehabilitation Research, Administration for Community Living, U.S. Department of Health and Human Services. (See <https://msktc.org/burn/model-system-centers> for more information).

Clinicians measure depth of burns in degrees:

- **First degree** burns are the mildest type of burn. They are also called superficial burns. First degree burns damage only the epidermis, which is the first layer of skin. First degree burns usually don't get infected or leave a scar. The skin may get red but won't break, and they heal within 3–5 days.
- **Second degree** burns are also called partial thickness burns. This type of burn damages the epidermis and the dermis (the second layer of skin). Second degree burns are painful. The injured area can swell and appear red with blisters. There are two subtypes of second degree burns: superficial and deep. Superficial second degree burns heal quicker (~2 weeks) and typically do not scar. Deep second degree burns take longer to heal and may require surgery depending on the size and location. They often have some degree of scarring.



- **Third degree** burns are also called full thickness burns. This type of burn goes through the epidermis and dermis and affects deeper tissues, which may also be damaged or destroyed. The injured area can appear charred and may be black, white, or deep red in color. This area is often numb to light touch. Third degree burns don't heal by themselves, so skin grafting is often necessary (discussed later).
- **Fourth degree** and deeper burns destroy the skin plus fat, muscle and sometimes bone.

How Is the Size of the Skin Burned Estimated?


Total body surface area (or TBSA)

burned is the percent (%) of the body that is burned. To estimate burn size, clinicians use a formula that considers the age of the injured person and a diagram called the Rule of Nines (shown below). This is the most common way to estimate burn size. First degree burns are not included in the calculation of percent TBSA burn.

How Is the Severity of the Burn Assessed?

Clinicians consider many factors to determine the severity of a burn injury:

- **Degree** (depth) of the burn,
- **Size** (percentage) of the skin that is burned, and
- **Age** of the injured person
- **Other factors** include:
 - Location of the burn on the body,
 - Inhalation of toxic gases or smoke,
 - Type of burn,
 - Other traumatic injuries (bone fractures), and
 - Other health issues (such as diabetes, heart problems, or alcoholism).



Can Lungs Be Burned?

When breathed in, smoke or toxic gases can harm the lungs. Clinicians call this an **inhalation injury**. This type of injury often occurs when the injured person is trapped in an enclosed area for a long amount of time. Damage depends on the type of gas and smoke particles inhaled and on length of exposure.



PHASE II: Healing the Burn

What Are Skin Grafts and Donor Sites?

Skin grafts are thin layers of skin that surgeons take from an unburned area and then surgically place on the burned area. The area where the skin is taken from is called the donor site. Common donor sites are the thigh and the back but may be taken from any uninjured area of the body other than the face. The donor site generally takes about two weeks to heal.



What Are the Different Types of Skin Grafts?

Surgeons consider many factors when deciding what type of skin graft to use. They look at the condition, thickness, and size of the wound and where the injury is on the body.

- **Autografts** are permanent skin grafts that replace burned skin. With this graft, surgeons remove skin from one place on the body and place it on the burned area of the body. There are two types of autografts:
 - **Split-thickness skin graft** involves removing the epidermis and a shallow layer of the dermis and then placing it on the burned area.
 - **Full-thickness skin graft** involves removing the epidermis and dermis and placing it on the burned area.
- **Allografts and homografts** are temporary grafts to cover the wound. Donor skin comes from another person (usually a cadaver).
- **Xenografts** are temporary grafts to cover the wound. Donor skin comes from a pig or fish.
- **Meshed grafts** are grafts in which donor skin is perforated with small slits or holes. Surgeons can then expand the mesh to cover a large burned area of the body. All four types of grafts can be meshed.
- **Sheet grafts** are grafts that are placed on the wound directly from the donor site without being meshed. These grafts are typically used for small surface area burns and in areas where function and cosmesis (cosmetic preservation) is important.

What Should Patients Expect?

The recovery process differs for everyone. Most patients report feeling pain, fatigue, and itching during recovery and rehabilitation.

- **Pain** is common. Third degree burns are painful with deep pressure. Second degree burns are painful with air movement or changes in temperature. First degree burns are painful on the surface of the skin. Pain can also occur in the process of dressing changes. Most importantly, a delayed onset of increased pain can be associated with onset of wound infection. Additional information can be found in the [“Managing Pain After Burn Injury”](#) factsheet.
- **Fatigue** is common. The more severe the burn injury, the greater the level of fatigue.
- **Itching** is a common part of the healing process. Patients should never scratch their wounds. Moisturizers and antihistamines can help make the burned area less itchy. Application of cool compresses on a healed burn may also help. Additional information can be found in the [“Itchy Skin After Burn Injury”](#) factsheet.



Patients may have a tough time dealing with these symptoms, but should always let their care provider know how they feel and see what options are available to help relieve symptoms. *Space is provided at the end of this factsheet for patients, family members and friends to write down questions.*



What About Infection?

Burned areas can get infected, at the time of the injury and during the healing process. To prevent infection, people with burn injuries should follow the treatment orders of their health care team. They should also follow the hospital's infection control guidelines, such as using gloves and gowns when recommended. Practicing good hand hygiene (clean hands) can help prevent infection. This applies to both the injured person and their family members, friends, and caregivers. It is also important that patients wash their wounds with mild soap and water at the time of each dressing change to decrease incidence of infection unless told otherwise. Dressings typically involve topical antibiotic ointment to prevent seeding of bacteria into the open wound bed.



What About Scarring?

Depending on the burn injury degree and location, along with age and ethnicity, scarring may occur. Unfortunately, it is difficult to predict who will develop scarring. Studies have shown that burns will typically not scar if they heal within 14 days. The longer a wound remains open, the greater the risk of scarring. Hypertrophic burn scars (raised scars in the area of the original burn) may occur and are a common complication in burn injuries. These scars may limit function depending on the location as well as affect body image. These scars may also become symptomatic and result in discomfort, itchiness, and tightness.

There are various treatments to manage hypertrophic scars including but not limited to pressure garments, silicone gel sheets, self-massage therapy and laser scar revision therapy. Consult with your provider for more information regarding these treatment modalities if you are interested.

PHASE III: Long-Term Recovery & Rehabilitation

What About Rehabilitation?

- **Rehabilitation** is the process of using education and therapy to help a patient reach his or her maximum potential for recovery after an injury. Rehabilitation may also include physical therapy or occupational therapy. Physical therapy treats physical impairments. It aims to improve a person's ability to walk and move, do daily tasks, and live independently. Treatment may include exercise, splinting, bandaging, medicine, and use of assistive devices (such as crutches or walkers), massage, and joint mobilization. Below are some of the common services provided as part of rehabilitation.
- **Occupational therapy** uses everyday activities and materials to help patients in their own recovery. For example, a patient could practice dressing, bathing, and feeding themselves or cooking.
- **Speech-language pathology** helps with swallowing, voice, cognitive issues, and communication. Several types of facilities provide rehabilitation services to patients with burn injuries.
- **A burn center** often provides intensive physical and occupational therapy during acute hospitalization. The burn team will also follow the patient long-term, prescribing additional therapy if needed.
- **An inpatient rehabilitation facility** is a specially designed medical rehabilitation facility. Patients stay at the facility full-time after discharge from the burn center and must be able to do at least 3 hours of therapy per day.
- **A skilled nursing facility** is a place where patients may go after they leave the burn center. Patients live in these facilities and receive medical and nursing care to help them recover.



- **An outpatient rehabilitation center** is a medical facility that provides physical and occupational therapy services to people who are living at home.

What About Nutritional Needs?

Eating well is a key part of the recovery process. Healing from a burn injury requires more calories and protein than healing from other types of injury. Additional information is provided in the [“Healthy Eating After Burn Injury”](#) factsheet. **Nutritionists** may be available to provide information about a well-rounded diet.



What About Mental Health?

It is common to feel anxious or upset after a traumatic injury. Burns can put tremendous stress on the patient and family. Burn teams include psychologists and psychiatrists who can help with feelings and concerns. **Psychiatric and psychological services** address issues like depression, anxiety, post-traumatic stress, addiction, and sleep problems. **Social services** can help families with managing housing, finances, and navigating the healthcare system. More information can be found in the [“Psychological Distress After Burn Injury”](#) factsheet.

What Can Family and Friends Do When a Loved One Is Injured and Hospitalized?

In addition to care from medical staff, patients can also draw support from family and friends. Visitors and caregivers play an important role during the recovery process. It is important for family members and friends to also take care of themselves (for example, housing, food, and care of children). It can be helpful to coordinate visiting times amongst friends and family members. Family members should also assign one person to be the main point of contact with medical staff. This person should be an adult who is available to talk with the burn team.

What Resources Are Available?

Factsheets from the Burn Model System are a great source of information and discuss many topics related to burn injury, such as sleep problems, body image, social interactions, and nutrition. Factsheets are available online (<http://www.msktc.org/burn/factsheets>) in English and Spanish.

Videos are also available online (<http://www.msktc.org/burn/videos>). They cover exercise and employment after a burn injury.

Feeling anxious and afraid is common after a traumatic injury. Several local and national support systems offer support and suggestions about recovering from a burn injury. Patients, family members, and friends should ask their health care team about options for support systems, such as the Phoenix Society for Burn Survivors (<https://www.phoenix-society.org/>). The Phoenix Society is a national **peer support** system in more than 60 hospitals and burn clinics throughout North America. This group is made up of burn survivors and their family members who are trained to offer support and can schedule a visit and share their recovery process.



Questions for the Burn Team

Use the space below to write down questions about the burn injury, treatment options, concerns about hospital care, and anything else to ask the health care team.

Reference

Gauglitz, G. (2023). Hypertrophic scarring and keloids following burn injuries. In M. Jeschke & K. Collins (Eds.), *UpToDate*. <https://www.uptodate.com/contents/hypertrophic-scarring-and-keloids-following-burn-injuries>



Authorship

Understanding a Burn Injury was originally developed in 2018 by Laura C. Simko, BS, and Emily A. Ohrtman, BA, from the Boston-Harvard Burn Model System; and Gretchen J. Carrougner, MN, RN, and Nicole S. Gibran, MD, FACS, from the Northwest Regional Burn Model System, in collaboration with the Model Systems Knowledge Translation Center (MSKTC). It was reviewed and updated in 2023 by John Schulz, MD, Lauren Shepler, MPH, and Lauren Tripodi, PA, in collaboration with the MSKTC.

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider about specific medical concerns or treatment. The contents of this factsheet were originally developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082) and were updated under NIDILRR grants (90DPKT0009 and 90DPBU0008). NIDILRR is a Center within the Administration for Community Living (ACL), U.S. Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, HHS, and you should not assume endorsement by the federal government.

Recommended citation: Schulz, J., Shepler, L., & Tripodi, L. (2023). *Understanding a burn injury* [Fact sheet]. Model Systems Knowledge Translation Center (MSKTC). <https://msktc.org/burn/factsheets/understanding-burn-injury>.

Copyright © 2023 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Understanding and Improving Body Image After Burn Injury

October 2023

<https://msktc.org/burn/factsheets>

BURN Factsheet

This fact sheet explains issues of importance for burn survivors concerning body image and practical strategies for use in social situations.

A burn injury, regardless of size, can change how the body looks and works. Burn injuries can also lead to experiencing body image distress for people with scars even if they are not readily visible. Body image refers to how happy, comfortable, and confident a person is with how they look. About one-third of burn survivors have severe distress about changes in the way their body looks, feels, and works when they are first hospitalized. Almost everyone has ups and downs as they heal but most children and adults get used to the change in their appearance over time.



What Causes Body Image Distress?

Many factors may cause body image distress after a burn injury. These include how a person feels about their burns; unhelpful coping strategies; and a person's gender, mental health history, pre-injury body image, and support network. Body image distress may involve:

- Grief or sadness about changes in appearance and physical abilities,
- Anxiety about social or intimate scenarios where scars may be seen,
- Anxiety about actual and expected questions and stares from people in the community,
- Worry about how people will react when they see the scars, and
- A desire to be with a trusted person when in public places.

These feelings are a common part of adjusting after a burn injury. The following sections describe ways you can ease this distress and feel more positive about your body.

Phases of Healing

After a burn injury, your skin goes through several phases of healing. During each phase, it's important to follow the advice of your burn team to improve how your skin heals. Doing things to help with healing can also improve confidence and feeling in control.

Wound Healing Phase

What you see: Lighter colored skin replaces open wounds. For people with darker skin, pink colored skin may replace the burn wound at first and then return to closer to uninjured skin color with time. Generally, the faster a wound heals, the less scarring will occur.



The Model Systems Knowledge Translation Center works with Burn Model System (BMS) centers to provide free research-based rehabilitation resources for people living with burn injury. (See <https://msktc.org/burn> for more information). This factsheet has been approved by experts from the BMS centers.



Model Systems
Knowledge Translation
Center

What you can do: Get involved with your wound care as much as possible, such as helping with the dressing changes. A cleaner wound heals faster. The more aware and involved you are with your wound care, the more likely you are to take good care of your burn.

Scar Formation Phase

What you see: After the wound heals, the skin changes over the next 3–4 months. It becomes darker, stiffer, and raised.

What you can do: Scars can cause distress because they look and feel different than uninjured skin. Scarring can also make skin stiff and painful.

- Work with your medical team to prevent and lessen scarring. Your team may tell you to wear pressure garments or splints, massage the scar, or do stretching exercises.
- Protect your healing skin from the sun by wearing protective clothing, like a hat or long sleeves, and applying sunscreen with SPF 30 or greater.



Scar Maturation Phase

What you see: Scar maturation can take up to 1–2 years. During this process, the scarred skin gradually returns to a more normal skin tone. It also becomes softer and flatter.

What you can do:

- The rehab team may tell you to keep wearing pressure garments or splints. The team may also tell you to keep massaging the scar and doing stretching exercises.
- Continue to protect your skin from the sun. For more information, visit: <https://msktc.org/burn/burn-topics/sun-protection-after-burn-injury>.
- You may want to meet with a surgeon who is trained in burn reconstruction. They may have tips for how to improve the appearance of scars and restore function. Some of their techniques may include lasers and cosmetic tattoos.

Burn injuries change how your skin looks. They can also change your appearance in other ways:

- Severe burns can damage structures under the skin. For example, when cartilage in the ears or nose is burned, there can be visible changes in these structures.
- Some burn survivors have many skin grafts and other reconstructive surgeries. These procedures can change the way a person looks.
- Healed skin or grafted skin may be permanently discolored. It may become lighter or darker than your uninjured skin.
- The appearance of burn scars may be difficult to change. If interested, ask trained make-up artists to assist with altering the appearance of your burn scars. For more information, visit: <https://resources.phoenix-society.org/makeup-application-for-burn-survivors>.
- Skin grafts can cause hair loss because hair follicles don't regrow.
- Sometimes burn injuries can result in the amputation or loss of fingers, toes, or limbs.



Wanting to look your best is normal and an important part of recovery. Even with the best care, rehabilitation, and reconstruction, burns can cause some permanent changes in how your body looks, feels, and works. Makeup,



clothing, or reconstructive surgery can help you feel more confident but part of the emotional healing process is learning to adjust to changes in your appearance. It may be helpful to focus less on your physical appearance and more on internal strengths and interests that make up your self-image. For example, focus on your accomplishments or roles that make you proud. These may include your education, career, or being a good friend or parent. Accepting your scars does not mean you have to like them. While some survivors report being completely comfortable in their changed bodies, others get their self-worth from internal qualities such as kindness and humor.



Here is an excerpt from a burn survivor talking about the change in their appearance:

"I don't *like* my appearance, some things not at all, but I *like* me. The person I am. The contributions I make. The impact I can have on others. The kindness I can show."

Social Interactions After Burn Injury

When seeing or meeting someone with burn scars for the first time, some people may stare, avoid contact, or ask pushy questions.

If This Happens, Here Are Some Things You Can Do:

- To feel confident when you're talking to someone, make eye contact. You should also use confident body language, smile, and use a friendly tone of voice.
- Have a quick response ready for when people ask "what happened," this can help relieve some anxiety when questions arise. Remember, you are in control of how much of your story you share! For example, "I was burned when I was younger, but fortunately I am back to doing all the activities I did before." Some burn survivors find that talking about their injury helps with emotional healing.
- If you don't want to discuss your appearance, you can say you don't want to talk about it. You also can guide the conversation to take the focus off you. Ask the other person open-ended questions (questions that a person can't answer with "yes" or "no"). For example, "I heard you went on a vacation. That sounds exciting. Tell me about it."
- When people stare or make negative comments, having a catch phrase—like "remember to be gracious"—can help you refocus and use social tools instead of reacting in a negative way.
- Some burn survivors find it helpful to be with a trusted person when in public places during the early stages of their recovery or return to the community.

Intimacy Following Burn Injury

You may be worried about showing your burn scars during intimate experiences. Here are some ways to help you feel more comfortable and confident:

- Talk to your partner. Learn about the different stages they may go through as you recover. These include withdrawal, avoidance, or being irritable with you.
- Encourage your partner to have contact with your skin. Moisturizing or massaging your scars can help both of you to overcome any hesitation or the "fear of rejection." Perhaps your partner can get familiar with the different feel and texture of your skin before you leave the hospital.
- Pursue grooming activities like styling your hair, showering, shaving, painting your nails, etc. This helps to improve your well-being and readjustments in body image.



- Get the support you need to address your concerns about intimacy. This may include talking to your health care team, other burn survivors, and/or organizations supportive of the burn community.

Other Resources That Can Help You With Social Interactions

- Social Interaction after Burn Injury (<https://msktc.org/burn/burn-topics/social-interaction-after-burn-injury>)
- Sexuality and Intimacy (<https://msktc.org/burn/burn-topics/sexuality-and-intimacy>)
- Gaining and Maintaining a Healthy Body Image from the Phoenix Society (<https://www.phoenix-society.org/resources/gaining-and-maintaining-a-healthy-body-image>)
- Changing Faces' Tools to Help You Cope with Other Peoples' Reactions (<https://www.changingfaces.org.uk/advice-guidance/coping-with-peoples-reactions/tools-cope-with-peoples-reactions/>)

Young Burn Survivors and Teasing

Parents and teachers need to closely watch how young burn survivors act and interact with other people. Children and teenagers often tease each other about even slight differences in appearance. Young burn survivors who are teased may become depressed or anxious when meeting new people. Children and teenagers often tease because they are curious or don't understand the situation or why their peers may look different.



Adults can intervene early to help protect kids from teasing and bullying through education. For example, it is helpful if adults explain the differences between curiosity (questions asked based on curiosity) versus teasing (malicious or mean intent). Young survivors may feel like they are being teased when their peers are simply being curious when asking about their scars. Second, adults can give their children and students, teachers, and coaches some basic information about burns and scarring. This might satisfy their curiosity in a less intrusive way. They can also teach others how to treat a burn survivor with respect. This can help create a supportive environment for young burn survivors. Schools often have policies in place to minimize teasing and bullying and create a supportive learning environment. Parents should review and communicate with school administration about these policies to decide how they can help their child.

School Reentry and Burn Camp Programs

Burn centers often have information and/or programs to help burn survivors return to school or may connect you with organizations supportive of the burn community that can help with this. Such programs may include inviting a burn expert to visit the burn survivor's school before the child returns. The burn expert explains the burn recovery process to the survivor's teachers and classmates. The burn expert also encourages students to be kind and supportive of the burn survivor. Many families find this process helpful in creating a supportive environment for the burn survivor.

Going Back to School After a Major Burn Injury and the Phoenix Society's *Return to School After a Burn Injury* has helpful information about the school reentry process. For more information on these resources, visit <https://msktc.org/burn/factsheets/going-back-school-after-major-burn-injury> and <https://www.phoenix-society.org/what-we-do/school-reentry>.



Some burn centers offer burn camps for children. These camps are often free. These camps give children a chance to play and interact with other children with burn injuries in order to build confidence and



practice social skills. Ask your burn center if they or other known organizations may offer these opportunities in your area.

Finding Help

Recovering from a burn can be tough emotionally. Get help if you feel anxious or depressed, have nightmares, or relive how you were injured. Many burn survivors find it helpful to talk to a mental health provider who has experience with the challenges of recovering from a burn. Your local burn center or health care provider can refer you to a mental health professional in your area.



You may want to talk to your health care provider about the following treatment options:

Cognitive behavioral therapy (CBT): CBT is a treatment approach used in behavioral health. CBT shows people how to understand and improve the connections between their thoughts, emotions, and behaviors. It is effective in treating severe depression and anxiety. CBT also helps people with body image distress.

Physical activity: Exercise may improve a person's body image. Exercise can increase energy and build confidence. It can also lessen feelings of hopelessness and lead to an appreciation of one's body. Even if you have never been physically active, you can start with 20 minutes of walking every day. Talk to your health care team about a structured exercise program.

Getting the Support You Need

With burn injuries, emotional healing is just as important as physical healing. Consider getting support from family, friends, colleagues, health professionals, and other burn survivors who may share your experience.

The Phoenix Society, a nonprofit organization based in the United States, offers the Phoenix Survivors Offering Assistance in Recovery (SOAR) program. This program connects people with new burn injuries to survivors and family members who have been impacted by a burn injury. Trained burn survivors are available across the United States to offer support. For more information, visit <https://www.phoenix-society.org/phoenix-soar>.

Additional Resources

The Model System Knowledge Translation Center website has many free research-based resources available, one that may be helpful for parents is "Help Your Child Recover—Build Your Child's Resilience after a Burn Injury." See: <http://www.msktc.org/burn/factsheets/Build-Childs-Resilience-After-Burn-Injury>.



The Phoenix Society is "dedicated to empowering anyone affected by a burn injury." For more information, visit <http://www.phoenix-society.org> or call 1-800-888-BURN. The Phoenix Society offers the following resources:

- Live, weekly chat about common issues for burn survivors and their loved ones (<https://www.phoenix-society.org/what-we-do/peer-support-chat>)
- Beyond Surviving: Social Skills online course (<https://resources.phoenix-society.org/beyond-surviving-whitepaper>)



Model Systems
Knowledge Translation
Center

Changing Faces is an advocacy organization in England. Its mission is “to create a better and fairer future for everyone who has a disfigurement to their face or body from any cause and their families.” For more information, visit <http://www.changingfaces.org.uk/Home>.

References

- Ajoudani, F., Jasemi, M., & Lotfi, M. (2018). Social participation, social support, and body image in the first year of rehabilitation in burn survivors: A longitudinal, three-wave cross-lagged panel analysis using structural equation modeling. *Burns*, 44(5), 1141–1150.
- Blakeney, P., Partridge, J., & Rumsey, N. (2007). Community integration. A review of the issues related to community integration of burn survivors. *Journal of Burn Care & Research*, 28(4), 598–601.
- Cleary, M., Kornhaber, R., Thapa, D., West, S., Visentin, D. (2020). A quantitative systematic review assessing the impact of burn injuries on body image. *Body Image*, 33, 47–65.
- Corry, N., Pruzinsky, T., & Rumsey, N. (2009). Quality of life and psychological adjustment to burn injury: social functioning, body image, and health policy perspectives. *International Review of Psychiatry*, 21(6), 539–548.
- Huang, Y. K., Su, Y. J. (2021). Burn severity and long-term psychosocial adjustment after burn injury: The mediating role of body image dissatisfaction. *Burns*, 47(6), 1373–1380.
- Kammerer Quayle, B. (2006). Behavioral skills and image enhancement training for burn survivors: essential interventions for improving quality of life and community integration. In R. Sood & B. Achauer (Eds.), *Achauer and Sood's burn surgery: reconstruction and rehabilitation*. Elsevier Health Sciences.
- King, I. (2018). Body image in paediatric burns: a review. (2018). *Burns & Trauma*. 6, 12.
- Partridge, J. (2006). From burns unit to board-room. *British Medical Journal*, 332(7547), 956–959.
- Thompson, A., & Kent, G. (2001). Adjusting to disfigurement: Processes involved in dealing with being visibly different. *Clinical Psychology Review*, 21(5), 663–682.
- Willemse, H., Geenen, R., Egberts, M. R., Engelhard, I. M., & Van Loey, N. E. (2023). Perceived stigmatization and fear of negative evaluation: Two distinct pathways to body image dissatisfaction and self-esteem in burn survivors. *Psychology & Health*, 38(4), 445–458.



Authorship

Understanding & Improving Body Image After Burn Injury was originally developed by John Lawrence, PhD, James Fauerbach, PhD, and Shawn Mason, PhD, in collaboration with Model Systems Knowledge Translation Center (MSKTC). It was reviewed and updated by Karen Kowalske, MD, Kimberly Roaten, PhD, Emma Turner, and Kyra Solis-Beach from the University of Texas Southwestern Medical Center and Parkland Burn Center/NTBRMS in collaboration with the MSKTC.

Source: The content in this factsheet is based on research and/or professional consensus. This content has been reviewed and approved by experts from the Burn Model System (BMS) centers, funded by the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR). The content of the factsheet has also been reviewed by individuals with burn injury and/or their family members.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this factsheet were developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082) and were updated under an NIDILRR grant (90DPKT0009 and 90DPBU0006). NIDILRR is a Center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this factsheet do not necessarily represent the policy of NIDILRR, ACL, or HHS, and you should not assume endorsement by the Federal Government.

Recommended citation: Kowalske, K., Roaten, K., Turner, E., & Solis-Beach, K. (2023). *Understanding and improving body image after burn injury*. Model Systems Knowledge Translation Center (MSKTC).

<https://msktc.org/burn/factsheets/understanding-and-improving-body-image-after-burn-injury>

Copyright © 2023 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



Model Systems
Knowledge Translation
Center

Wound Care After Burn Injury

September 2017

www.msktc.org/burn/factsheets

BURN Factsheet

This factsheet explains how to take care of a burn injury. It gives an overview of the three degrees of burns and the standard treatments for each. It also discusses the types of products available to treat burn injuries. The wound care described here is following discharge from the hospital.

The Burn Model System is sponsored by the National Institute of Disability, Independent Living, and Rehabilitation Research, U.S. Department of Health and Human Services' Administration for Community Living. (See <http://www.msktc.org/burn/model-system-centers> for more information).

Understanding the Extent of Your Burn


Burn injuries are caused by fires or flames, hot liquids or steam, contact with a hot object or agent like grease or tar, chemicals, or electricity. When evaluating a burn injury, doctors look at two factors: how deep the burn is and the burn size which is measured by the *percent total body surface area (% TBSA)*. The burn depth depends on how hot the agent was and how long the burned area was in contact with the agent and how thick the skin is in the area. There are three levels of a burn injury:

- *First-degree burns* affect the top layer of skin, called the epidermis (ep-i-DUR-mis). These burns cause minor damage to the skin. Skin may be red and tender or swollen. An example would be a mild sunburn that turns red and may peel. First-degree burns can generally be treated at home.
- *Second-degree burns* (also called partial thickness burns) go through the second layer of skin, called the dermis (DUR-mis). These burns cause pain, redness, and blisters and are often painful. The injury may ooze or bleed. They usually heal within 1 to 3 weeks. After healing, skin may be discolored. These burns generally do not leave raised scars. Treatment for second-degree burns varies. It may include ointments or special dressings. Surgery may be necessary for very deep second degree burns or those that are slow to heal.
- *Third-degree and more severe burns* (also called full thickness burns) damage both layers of the skin and may also damage the underlying bones, muscles, and tendons. Injured skin may turn white, black, and/or gray. It may feel dry and leathery. Sometimes there is no pain because the nerve endings under the skin are destroyed. Third-degree burns have a high risk of infection. They are usually treated with skin grafts. This surgery, done with general anesthesia, removes the injured skin and replaces it with healthy skin from an uninjured area of the body. Full thickness burns that are not grafted may take months or even years to heal. Third-degree burns likely leave raised scars. Burn survivors may have a combination of first, second, and third degree burns. Talk with your health care providers to better understand your specific injuries.

Treatment Options for Burn Injury

Wound Care

Antibiotic (an-ti-bahy-OT-ik) ointments or creams are often used to prevent or treat infections in patients with second-degree burns. Using these ointments may require the use of bandages. Dressings may need to be changed daily. This can be a painful process. Your doctor can assist you in coordinating the dressing changes with your pain medication. Dressings can be soaked off with water in a sink or shower. The skin and the burn wound should be washed gently with mild soap and rinsed well with tap water. Use a soft wash cloth or piece of gauze to gently remove old medications. A small amount of bleeding is common with dressing changes. Your doctor will decide on the appropriate dressing and ointment. This will be based on the location of the burn, the need to control drainage, and your comfort.



There are many “advanced wound care products” available for burns. These products don’t require daily dressing changes and can be left in place until the wound heals. This can make pain control much easier and may decrease anxiety about wound care. These types of dressings include impregnated (im-PREG-neyt-ed) gauzes, foams, honey, and silver dressings. Many of the currently available dressings are combinations of these categories. There are many different brand names. Your burn care team will determine the most appropriate product to use. They will also decide when to apply and remove it.

Skin Grafts

Larger areas of third degree (full thickness) burns are treated with skin grafts. This surgery removes dead skin and replaces it with healthy skin from another part of the body. The grafted skin is often treated with an antibiotic ointment and a nonstick dressing. There are three types of skin grafts.

- *Sheet grafts* are usually applied to the face or hands for better cosmetic effect. Sheet graft uses the whole piece of skin without the holes in it. It gives a better cosmetic appearance but requires much more skin to cover a specific area. Newly healed grafts are very fragile. Special care should be taken to protect them. Be careful not to bump, rub, or scratch them. Do not wear rough clothing or anything that rubs; this can cause blistering.
- *Meshed grafts* are used for larger wounds. For permanent wound coverage, a piece of your own skin is taken from another part of the body (donor skin) to close the open area. When the donor skin is taken off the body, it shrinks. To stretch the donor skin, it is put through a machine that makes small slits or holes in the skin. This stretched skin covers a larger area than an unmeshed sheet graft, but leaves a permanent mesh pattern similar to stockings. The wound heals as the areas between the meshed graft and the holes fill in with new skin. Once the mesh sheet sticks to the skin and the drainage stops, the wound is considered healed and can be left open to air. Lotion can be used to keep it moist.
- *Full-thickness grafts* are used for reconstruction of small areas that are prone to contracture such as the hand or chin. It consists of the full thickness of the skin and shrink the least compared to other grafts.

The area of the donor site is similar to a second-degree burn. Most burn providers use one of the advanced wound dressings that can be left in place for 7–14 days while healing occurs. Any remaining small open areas on the donor site can be treated with antibiotic ointment. Notify your burn provider of any areas of redness, warmth, and increased pain. These can be symptoms of an infection.

Moisturizing

Moisturizing will be very important after burn injury.

- Once the skin is closed and no longer draining, it is important to keep it well moisturized. This decreases the chances of developing blisters or skin tears. It also decreases itching and can make movement easier.
- There are many different lotions available.
- Lotions in bottles have a higher water content. They are also easier to apply. They often need to be applied frequently.
- Lotions in tubes and jars are thicker. They need to be massaged in more thoroughly. They last longer on your skin.
- Unscented lotion should be applied and massaged into the scar several times per day. Unscented lotion is important – ask your doctor for recommendations.
- Applying lotion is a good time to touch your scars using light pressure; touching your scars is helpful in keeping them from getting sensitive. It is also a good time to do a little stretching. This factsheet has more information about stretching:
<http://www.msktc.org/burn/factsheets/Exercise-After-Burn-Injury>.

Blisters

- Newly healed skin is fragile. Minor shearing (rubbing force) on the scar can cause blisters. Blisters also can develop from clothes that fit too tightly, shearing while putting on pressure garments, or rubbing or scratching the burn scar.
- Blisters should be pierced and drained as soon as you notice them. Use a sterile (STER-il) needle to make a small hole. Then drain the blister onto a piece of gauze. Put a little antibiotic ointment on the area.
- If a blister opens up, you might need to bandage it with a nonstick dressing. Do not use adhesive or sticky bandages or tape that is difficult to take off. Your skin might tear.
- Over moisturizing especially on face can occlude pores and cause pimples. If this occurs, reduce moisturizing and consult your doctor or the burn team.

Skin Tears

- Skin tears occur when you bump into something such as a doorway, a counter top, or a piece of furniture. They can also be caused by scratching.
- If the area bleeds, put firm pressure over the wound for about 5 minutes until the bleeding stops.
- Wash the area gently and thoroughly with mild soap and water.
- Use a small amount of antibiotic ointment and a nonstick dressing and allow the wound to heal. If the surrounding area becomes red and warm, you might have an infection. Contact your health care provider for further evaluation.
- If the wound continues to crack open, get bigger, or deepen, your health care provider may recommend a splint or cast. This will keep the wound area still.

Ulcerations

- Ulcerations (uhl-suh-REY-shuhns) are breakdowns in the skin. They usually occur across bands of scar tissue around your shoulder, the front of your elbow, and the back of your knee.
- These areas can be difficult to heal. Physical movements like exercise can cause the wound to continuously crack open or get bigger.
- Keep the wound covered with a thin film of antibiotic ointment. Keep the surrounding skin well moisturized, especially when you are exercising and stretching.

Allergic skin reactions

- Be sure to let your doctor know if you have any skin allergies.
- Allergic skin reactions can be caused by using antibiotic ointments for a long time. Other causes include changing the type of lotion or soap that you use, changing laundry detergents, or changing the elastic in pressure garments.
- If you have an allergic skin reaction, stop using all soaps, lotions, and ointments for 2 to 3 days.
- Once the reaction has gone away, you can start using soaps and moisturizers again. Add these products back into your daily routine one at a time.
- Allow 2 to 3 days in between adding each product back into your daily routine. This will allow time to see if the reaction returns.

What Can You Do?

- Be active in your recovery. Ask questions and help make decisions about your care.
- Take a list of questions or concerns to your medical appointments for your health care provider to address.
- Follow your provider's wound care instructions.
- Always keep your skin clean and well moisturized.
- Avoid bumping, scraping, or scratching.
- Start exercises as soon as possible. Refer to this factsheet for more information about exercise after your burn:
<http://www.msktc.org/burn/factsheets/Exercise-After-Burn-Injury>

Additional Resources

For more information regarding the care of your wounds, please contact your doctor or therapist so that they can address your specific needs.

Johnson, R.M., and Richard, R. (2003). "Partial-Thickness Burns: Identification and Management." *Advances in Skin Wound Care*, 16 (4), 178-287.

Kowalske, K. (2011). "Burn Wound Care." *Physical Medicine & Rehabilitation Clinics of North America*, 22, 213-227.

Honari, S. (2004). "Topical Therapies and Antimicrobials in the Management of Burn Wounds." *Critical Care Nursing Clinics of North America*, 16 (1), 1-11.



Authorship

Wound Care After Burn Injury was developed by Karen J. Kowalske, MD, Sandra Hall, PT, DPT., Radha Holavanahalli, PhD, and Lynne Friedlander, M.Ed, in collaboration with the Model Systems Knowledge Translation Center.

Source: Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the Burn Injury Model Systems.

Disclaimer: This information is not meant to replace the advice of a medical professional. You should consult your health care provider regarding specific medical concerns or treatment. The contents of this factsheet were developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90DP0082). The contents of this factsheet do not necessarily represent the policy of Department of Health and Human Services, and you should not assume endorsement by the Federal Government.

Copyright © 2017 Model Systems Knowledge Translation Center (MSKTC). May be reproduced and distributed freely with appropriate attribution. Prior permission must be obtained for inclusion in fee-based materials.



